

THE FARRELL-HSIANG METHOD REVISITED

BARTELS, A. AND LÜCK, W.

ABSTRACT. We present a sufficient condition for groups to satisfy the Farrell-Jones Conjecture in algebraic K -theory and L -theory. The condition is formulated in terms of finite quotients of the group in question and is motivated by work of Farrell-Hsiang.

INTRODUCTION

Farrell-Hsiang used in [7] a beautiful combination of controlled topology and Frobenius induction to prove that the Whitehead group of fundamental groups of compact flat Riemannian manifolds is trivial. This general method has been refined and used further by Farrell-Hsiang, Farrell-Jones and Quinn, see for example [8, 9, 11, 17]. These results belong to a much wider collection of results that ultimately led to the Farrell-Jones Conjecture in [12] that predicts a largely homological formula for K - and L -theory of group rings. For more information about the Farrell-Jones Conjecture and its applications we refer to [5, 15].

The present paper gives an axiomatic treatment of the Farrell-Hsiang method leading us to the definition of Farrell-Hsiang groups below. More generally we define a group to be a Farrell-Hsiang group with respect to a given family of subgroups \mathcal{F} , more or less if the Farrell-Hsiang method is applicable relative to \mathcal{F} . Our main result states that Farrell-Jones Conjecture holds for these groups and this family. In the most important case \mathcal{F} is the family VCyc of virtually cyclic subgroups. In this case our result implies that if a group G is a Farrell-Hsiang group relative to VCyc, then G satisfies both the K - and L -theoretic Farrell-Jones Conjecture with coefficients in additive categories. This result is used in work with Tom Farrell to prove the Farrell-Jones Conjecture for poly-cyclic groups [1].

In [10] Farrell-Jones used a wonderful combination of controlled topology and the dynamics of the geodesic flow on negatively curved manifolds to prove that the Whitehead group of the fundamental group of such manifolds vanishes. This Farrell-Jones method has also been refined and further used in many papers about the Farrell-Jones conjecture and the Borel conjecture, see for example [12, 13]. In [4, 2] an axiomatic treatment for this method is given that is from a formal point of view very similar to our treatment here. In both cases a transfer and a contracting map are the main ingredients. The main difference is, that the transfer in the Farrell-Hsiang method uses a finite discrete fiber and its construction depends on Frobenius induction, whereas in the Farrell-Jones method the fiber is a compact contractible space and the transfer is essentially given by the tensor product with the singular chain complex of this fiber. Also, in applications the construction of the contracting maps is very different. In the first case subgroups of finite but large index are exploited, in the second case the dynamic of flow spaces is a key ingredient.

Date: January 2011.

1991 Mathematics Subject Classification. 18F25, 19A31, 19B28, 19G24.

Key words and phrases. Farrell-Jones Conjecture, K - and L -theory of group rings.

Acknowledgements. The first author thanks Frank Quinn for a long email exchange about the Farrell-Hsiang method. This paper was supported by the SFB 878 *Groups, Geometry and Actions* and by the Leibniz-award of the second author.

1. FARRELL-HSIANG GROUPS

A finite group H is said to *hyperclementary* if it can be written as an extension $1 \rightarrow C \rightarrow H \rightarrow P \rightarrow 1$, where C is a cyclic group and P is a p -group for some prime p .

Definition 1.1 (Farrell-Hsiang group). Let \mathcal{F} be a family of subgroups of the finitely generated group G . We call G a *Farrell-Hsiang group* with respect to the family \mathcal{F} if the following holds for a fixed word metric d_G :

There exists a natural number N such that for every natural number n there is a surjective homomorphism $\alpha_n: G \rightarrow F_n$ with F_n a finite group such that the following condition is satisfied. For any hyperclementary subgroup H of F_n we set $\overline{H} := \alpha_n^{-1}(H)$ and require that there exists a simplicial complex E_H of dimension at most N with a cell preserving simplicial \overline{H} -action whose stabilizers belong to \mathcal{F} , and an \overline{H} -equivariant map $f_H: G \rightarrow E_H$ such that $d_G(g, h) < n$ implies $d_{E_H}^1(f(g), f(h)) < \frac{1}{n}$ for all $g, h \in G$, where $d_{E_H}^1$ is the l^1 -metric on E_H .

Theorem 1.2 (Main Theorem). *Let G be a Farrell-Hsiang group with respect to the family \mathcal{F} in the sense of Definition 1.1. Then G satisfies the K -theoretic and L -theoretic Farrell-Jones Conjecture with additive categories as coefficients with respect to the family \mathcal{F} .*

For the precise formulation and discussion of the Farrell-Jones Conjecture with coefficients in additive categories we refer to [3].

Remark 1.3. Definition 1.1 can be weakened if one is only interested in the L -theoretic Farrell-Jones conjecture. In this case it suffices to consider all subgroups H of F that are either 2-hyperclementary or p -elementary for some prime $p \neq 2$. In other words p -hyperclementary subgroups that are not p -elementary can be ignored for all odd primes p .

2. CATEGORICAL PRELIMINARIES

Additive G -categories with involutions. In this paper we will understand notions like additive category (with involution) or additive G -category (with involution) always in the strict sense. This means that all our additive categories will come with a strictly associative functorial direct sum $(M, N) \mapsto M \oplus N$ and an involution I on an additive category \mathcal{B} is a contravariant functor $I: \mathcal{B} \rightarrow \mathcal{B}$ with $I^2 = \text{id}_{\mathcal{B}}$. When we talk about an additive G -category, the (right) G -action is understood to be in the strict sense, i.e., for every $g \in G$ we have a functor $R_g: \mathcal{B} \rightarrow \mathcal{B}$ of additive categories such that $R_h \circ R_g = R_{gh}$ for $g, h \in G$. If \mathcal{B} comes with an involution $I_{\mathcal{B}}$, then we require $I_{\mathcal{B}} \circ R_g = R_g \circ I_{\mathcal{B}}$ for all $g \in G$.

Remark 2.1. Often a more general definition of additive categories with involutions is used, where the equality $I^2 = \text{id}_{\mathcal{B}}$ is replaced by a natural equivalence $E: I^2 \rightarrow \text{id}_{\mathcal{B}}$. One may also consider additive categories with weak G -actions. We refer to [3], where all these notions are explained, it is shown how one can replace the weak versions by equivalent strict versions, and – most important – that for a proof of the Farrell-Jones Conjecture it suffices to consider the strict versions (see [3, Theorem 0.2]). We use the strict versions to simplify some formulas. The only slight disadvantage of this is, that it forces us to replace some very natural categories by some slightly less natural categories, see for instance the definition of $\text{mod}_{\mathbb{Z}}$ below.

A functor between additive categories with involutions (\mathcal{B}, I) and (\mathcal{B}', I') is a pair (F, E) where $F: \mathcal{B} \rightarrow \mathcal{B}'$ is an additive functor, and $E: F \circ I \rightarrow I' \circ F$ is a natural equivalence such that $I'(E(M)) = E(I(M))$ for all objects $M \in \mathcal{B}$. If $F \circ I = I' \circ F$ and $E = \text{id}$, then the functor is said to be strict. Most of our functors will be strict, but not all of them. Functors between additive categories with involutions induce maps in L -theory.

The category $\text{mod}_{\mathbb{Z}}$ of based finitely generated free abelian groups. On the category of finitely generated abelian groups the involution $T \mapsto T^* := \text{Hom}_{\mathbb{Z}}(-, \mathbb{Z})$ is not strict since T is not $(T^*)^*$ on the nose. To fix this inconvenience we will consider the following additive category with involution $\text{mod}_{\mathbb{Z}}$ instead. The objects of $\text{mod}_{\mathbb{Z}}$ are \mathbb{Z}^n , $n = 0, 1, 2, \dots$. The set of morphisms $\text{mor}_{\text{mod}_{\mathbb{Z}}}(\mathbb{Z}^n, \mathbb{Z}^m)$ is given by $n \times m$ -matrices. Composition is given by the usual matrix multiplication. The direct sum is given by $\mathbb{Z}^n \oplus \mathbb{Z}^m = \mathbb{Z}^{n+m}$. The involution on $\text{mod}_{\mathbb{Z}}$ acts as the identity on objects and as transposition of matrices on morphisms. For an additive category \mathcal{A} there is a functor

$$-\otimes_{\mathbb{Z}} -: \text{mod}_{\mathbb{Z}} \times \mathcal{A} \rightarrow \mathcal{A}$$

defined by $\mathbb{Z}^n \otimes_{\mathbb{Z}} M = \bigoplus_{i=1}^n M$, see for example [2, Section 6]. This functor is bilinear on morphisms groups. It follows that given an object \mathbb{Z}^n in $\text{mod}_{\mathbb{Z}}$, the functor $\mathbb{Z}^n \otimes_{\mathbb{Z}} -: \mathcal{A} \rightarrow \mathcal{A}$ is a functor of additive categories, and given an object $M \in \mathcal{A}$, the functor $-\otimes_{\mathbb{Z}} M: \text{mod}_{\mathbb{Z}} \rightarrow \text{mod}_{\mathbb{Z}}$ is a functor of additive categories. If \mathcal{A} comes with an involution, then $\text{mod}_{\mathbb{Z}} \times \mathcal{A}$ inherits the obvious product involution and $-\otimes_{\mathbb{Z}} -$ is compatible with the involutions.

The category $\text{mod}_{(\mathbb{Z}, G)}$ of $\mathbb{Z}G$ -modules which are finitely generated free as abelian groups. Let G be a group. We define the following additive category with involution $\text{mod}_{(\mathbb{Z}, G)}$. Objects in $\text{mod}_{(\mathbb{Z}, G)}$ are pairs (\mathbb{Z}^n, ρ) where $\rho: G \rightarrow \text{GL}(n\mathbb{Z})$ is a group homomorphism. A morphism $f: (\mathbb{Z}^n, \rho) \rightarrow (\mathbb{Z}^m, \eta)$ is a morphism $f: \mathbb{Z}^n \rightarrow \mathbb{Z}^m$ in $\text{mod}_{\mathbb{Z}}$ which is compatible with the homomorphisms ρ and η , i.e., $\eta(g) \circ f = f \circ \rho(g)$ for all $g \in G$. The direct sum is given by the direct sum in $\text{mod}_{\mathbb{Z}}$. Define an involution $I_{\text{mod}_{(\mathbb{Z}, G)}}$ on $\text{mod}_{(\mathbb{Z}, G)}$ as follows. It sends an object (\mathbb{Z}^n, ρ) to the object (\mathbb{Z}^n, ρ^*) , where $\rho^*(g)$ is defined by $I_{\text{mod}_{\mathbb{Z}}}(\rho(g^{-1}))$. A morphism $f: (\mathbb{Z}^n, \rho) \rightarrow (\mathbb{Z}^m, \eta)$ is sent to the morphism given by $I_{\text{mod}_{\mathbb{Z}}}(f)$.

Of course $\text{mod}_{(\mathbb{Z}, G)}$ is a model for the category of $\mathbb{Z}G$ -modules which are finitely generated free as abelian groups and has the extra feature that the involution is strict.

Let $\alpha: H \rightarrow G$ be a group homomorphism. We obtain a functor of additive categories with involution called *restriction*

$$\text{res}_{\alpha}: \text{mod}_{(\mathbb{Z}, G)} \rightarrow \text{mod}_{(\mathbb{Z}, H)},$$

which sends an object (\mathbb{Z}^n, ρ) to the object $(\mathbb{Z}^n, \rho \circ \alpha)$ and a morphism $f: (\mathbb{Z}^n, \rho) \rightarrow (\mathbb{Z}^m, \eta)$ to the morphism $f: (\mathbb{Z}^n, \rho \circ \alpha) \rightarrow (\mathbb{Z}^m, \eta \circ \alpha)$.

Next we define the induction functor for a subgroup H of G of finite index

$$\text{ind}_H^G: \text{mod}_{(\mathbb{Z}, H)} \rightarrow \text{mod}_{(\mathbb{Z}, G)}.$$

It will depend on a choice of representatives $g_0, \dots, g_{m-1} \in G$ for G/H . This choice will not matter in the sequel, since for two such choices we obtain a unique natural equivalence of the corresponding functors of additive categories with involution. Consider an object (\mathbb{Z}^n, ρ) in $\text{mod}_{(\mathbb{Z}, H)}$. The image under ind_H^G is the object $(\mathbb{Z}^{m \cdot n}, \eta)$, where $\eta(g) \in \text{GL}(m \cdot n, \mathbb{Z})$ for $g \in G$ is the morphism in $\text{mod}_{\mathbb{Z}}$ given by the matrix whose entry at $(kn + i, k'n + i')$ is 0 if $gg_k H \neq g_{k'} H$, and is $\rho(g_{k'}^{-1} g g_k)_{i, i'}$ if $gg_k H = g_{k'} H$. Here $0 \leq k, k' \leq m-1$, $1 \leq i, i' \leq n$.

Remark 2.2. Our above definition of ind_H^G may appear unnatural. But the only reason for this is our choice of the category $\text{mod}_{(\mathbb{Z}, H)}$; it really is the usual definition of induction:

Let (\mathbb{Z}^n, ρ) be an object of $\text{mod}_{(\mathbb{Z}, H)}$. Then \mathbb{Z}^n becomes an $\mathbb{Z}[H]$ -module via ρ . We have the following isomorphism of \mathbb{Z} -modules

$$\mathbb{Z}[G] \otimes_{\mathbb{Z}[H]} \mathbb{Z}^n \cong \bigoplus_{j=0}^{m-1} \mathbb{Z}[g_j H] \otimes_{\mathbb{Z}[H]} \mathbb{Z}^n \cong \bigoplus_{j=0}^{m-1} \mathbb{Z}^n \cong \mathbb{Z}^{m \cdot n}$$

and the above formula for η describes how the action of G on $\mathbb{Z}[G] \otimes_{\mathbb{Z}[H]} \mathbb{Z}^n$ conjugates to an action on $\mathbb{Z}^{m \cdot n}$ under the above isomorphism.

3. THE OBSTRUCTION CATEGORY $\mathcal{O}^G(E, Z, d; \mathcal{A})$

Let E be a G -space and (Z, d) be a quasi-metric space with a free, proper and isometric G -action. In this section we will review the additive category $\mathcal{O}^G(E, Z, d; \mathcal{A})$ that was originally defined in [4, Section 3], see also [2, Section 4]. If \mathcal{A} is an additive category with involution, then $\mathcal{O}^G(E, Z, d; \mathcal{A})$ is an additive category with involution.

Objects. Objects in $\mathcal{O}^G(E, Z, d; \mathcal{A})$ are given by sequences $M = (M_y)_{y \in Z \times E \times [1, \infty)}$ of objects from \mathcal{A} subject to the following conditions.

- (i) *G -compact support over $Z \times E$.* There is a compact subset K of $Z \times E$ such that $M_{z, e, t} = 0$ whenever $(z, e) \notin G \cdot K$.
- (ii) *Locally finiteness.* For all $y \in Z \times E \times [1, \infty)$ there existence an open neighborhood U such that $\{y \in U \mid M_y \neq 0\}$ is finite.
- (iii) *G -equivariance.* For all $y \in Z \times E \times [1, \infty)$ and $g \in G$ we have $M_{gy} = g(M_y)$. Here $gy = (gz, ge, t)$ for $y = (z, e, t)$.

The involution $I_{\mathcal{O}}$ on $\mathcal{O}^G(E, Z, d; \mathcal{A})$ acts on objects point-wise, i.e., we have $(I_{\mathcal{O}}(M))_{z, e, t} = I_{\mathcal{A}}(M_{z, e, t})$.

Morphisms. Let $M = (M_y)_{y \in Z \times E \times [1, \infty)}$, $N = (N_y)_{y \in Z \times E \times [1, \infty)}$ be objects from $\mathcal{O}^G(E, Z, d)$. A morphism $\psi: M \rightarrow N$ in $\mathcal{O}^G(E, Z, d)$ is given by a sequence $\psi = (\psi_{y, y'})_{y, y' \in Z \times E \times [1, \infty)}$ of morphisms $\psi_{y, y'}: M_{y'} \rightarrow N_y$ in \mathcal{A} subject to the following conditions.

- (i) *Row and column finiteness.* For all $y \in Z \times E \times [1, \infty)$ the set $\{y' \mid \psi_{y, y'} \neq 0 \text{ or } \psi_{y', y} \neq 0\}$ is finite.
- (ii) *Metric control over Z .* There is $R > 0$ (depending on ψ) such that $\psi_{y, y'} = 0$ whenever $y = (z, e, t)$, $y' = (z', e', t')$ with $d(z, z') > R$.
- (iii) *Metric control over $[1, \infty)$.* There is $A > 0$ (depending on ψ) such that $\psi_{y, y'} = 0$ whenever $y = (z, e, t)$, $y' = (z', e', t')$ with $|t - t'| > A$.
- (iv) *G -continuous control over $E \times [1, \infty)$.* Let $e_0 \in E$, V be an G_{e_0} -invariant neighborhood of e_0 and $b > 0$. (Here $G_{e_0} = \{g \mid ge_0 = e_0\}$.) Then we require the existence of $B > 0$ and a G_{e_0} -invariant neighborhood U of e_0 such that $\psi_{y, y'} = \psi_{y', y} = 0$ whenever $y = (z, e, t)$, $y' = (z', e', t')$ with $(e, t) \in U \times (B, \infty)$ and $(e', t') \notin V \times (b, \infty)$.
- (v) *G -equivariance.* For all $y, y' \in Z \times E \times [1, \infty)$ and $g \in G$ we have $\psi_{gy, gy'} = g(\psi_{y, y'})$.

For the constructions in this paper the second condition will be the most important condition and we will say that ψ is *R -controlled* if it is satisfied for a given $R > 0$. Addition and composition of morphisms is defined as for matrices: $(\psi + \psi')_{y, y'} = \psi_{y, y'} + \psi'_{y, y'}$ and $(\psi \circ \psi')_{y, y''} = \sum_{y'} \psi_{y, y'} \circ \psi_{y', y''}$. The involution is on morphisms defined by the formula $(I_{\mathcal{O}}(\psi))_{y, y'} = I_{\mathcal{A}}(\psi_{y', y})$.

We will often drop \mathcal{A} from the notation and write $\mathcal{A}^G(E, Z, d)$ instead of $\mathcal{A}^G(E, Z, d; \mathcal{A})$.

Functoriality. In this paper we will only need the functoriality of $\mathcal{O}^G(E, Z, d; \mathcal{A})$ in the Z -variable. Let (Z, d) and (Z', d') be quasi-metric spaces with free, proper and isometric G -actions. Let $f: Z \rightarrow Z'$ be G -equivariant continuous map such that for any $r > 0$ there is $R > 0$ such that $d'(f(z_0), f(z_1)) < R$ whenever $d(z_0, z_1) < r$. Then f induces a functor $f_*: \mathcal{O}^G(E, Z, d) \rightarrow \mathcal{O}^G(E, Z', d')$ which is given by $(f_*(M))_{e,z,t} = \bigoplus_{z' \in f^{-1}(z)} M_{e,z',t}$. (The condition ensures that metric control over Z is turned into metric control over Z' ; the G -compact support condition for objects ensures that the sum in the definition of f_* is finite.) Strictly speaking f_* is only defined up to natural equivalence because the direct sum may only be defined up to canonical isomorphism. (By our assumptions on \mathcal{A} only imply that sums over ordered finite index set are canonically defined.)

$\mathcal{O}^G(E, G, d)$ as the obstruction to the Farrell-Jones conjecture. The following result is a consequence of [2, Theorem 5.2].

Theorem 3.1. *Let G be a finitely generated group, d_G a word metric on G and \mathcal{F} be a family of subgroups.*

- (i) *Assume that $K_*(\mathcal{O}^G(E_{\mathcal{F}}G, G, d_G))$ is trivial in all degrees. Then the K -theory assembly map $H_*^G(E_{\mathcal{F}}G; \mathbf{K}_{\mathcal{A}}) \rightarrow K_*(\int_G \mathcal{A})$ is an isomorphism.*
- (ii) *Assume that $L_*(\mathcal{O}^G(E_{\mathcal{F}}G, G, d_G))$ is trivial in all degrees. Then the L -theory assembly map $H_*^G(E_{\mathcal{F}}G; \mathbf{L}_{\mathcal{A}}) \rightarrow L_*(\int_G \mathcal{A})$ is an isomorphism.*

The controlled product category. Let (Z_n, d_n) be a sequence of quasi-metric spaces with free, proper and isometric G -actions. Consider the product category $\prod_{n \in \mathbb{N}} \mathcal{O}^G(E, Z_n, d_n)$. A morphism $\varphi = (\varphi_n)_{n \in \mathbb{N}}$ is said to be R -controlled for $R > 0$ if φ_n is R -controlled for all n . We define $\mathcal{O}^G(E, (Z_n, d_n)_{n \in \mathbb{N}})$ as the category whose objects are objects from the product category and whose morphisms are morphisms from the product category that are R -controlled for some R . There is for any k a canonical projection functor $\mathcal{O}^G(E, (Z_n, d_n)_{n \in \mathbb{N}}) \rightarrow \mathcal{O}^G(E, Z_k, d_k)$.

4. THE CORE OF THE PROOF OF THE MAIN THEOREM 1.2

Let G be a Farrell-Hsiang group with respect to \mathcal{F} . Let N be the number appearing in Definition 1.1. For $n \in \mathbb{N}$ there is then $\alpha_n: G \rightarrow F_n$, a surjective group homomorphism onto a finite group F_n , such that the following holds: For any hyper-elementary subgroup H of F_n and $\bar{H} := \alpha_n^{-1}(H)$ there is a simplicial complex E_H of dimension at most N with a cell preserving simplicial \bar{H} -action whose stabilizers belong to \mathcal{F} , and an \bar{H} -equivariant map $f_H: G \rightarrow E_H$ such that $d_G(g, h) < n$ implies $d_{E_H}^1(f_H(g), f_H(h)) < \frac{1}{n}$ for all $g, h \in G$, where $d_{E_H}^1$ is the l^1 -metric on E . Here we write \bar{H} for $\alpha_n^{-1}(H)$ and we will use this convention throughout the remainder of this paper. We denote by \mathcal{H}_n the family of hyper-elementary subgroups of F_n . We set $X_n := G \times \prod_{H \in \mathcal{H}_n} \text{ind}_{\bar{H}}^G E_H$ and $S_n := G \times \prod_{H \in \mathcal{H}_n} G/\bar{H}$. We equip X_n and S_n with diagonal G action. We will use the quasi-metrics d_{X_n} on X_n and d_{S_n} on S_n defined by

$$d_{X_n}((g, x), (h, y)) := d_G(g, h) + n \cdot d_{\text{ind}_{\bar{H}}^G E_H}^1(x, y),$$

$$d_{S_n}((g, a\bar{H}), (h, b\bar{K})) := \begin{cases} d_G(g, h) & \text{if } K = H \text{ and } a\bar{H} = b\bar{K}, \\ \infty & \text{otherwise.} \end{cases}$$

Here $g, h, a, b \in G$, $x, y \in X_n$, $H, K \in \mathcal{H}_n$ and $d_{\text{ind}_{\bar{H}}^G E_H}^1$ is the l^1 -metric on $\text{ind}_{\bar{H}}^G E_H$. Abbreviate $E := E_{\mathcal{F}}G$. The proof of Theorem 1.2 is organized around the following

diagram of additive categories and functors.

$$(4.1) \quad \begin{array}{ccc} & & \oplus_{n \in \mathbb{N}} \mathcal{O}^G(E, X_n, d_{X_n}) \\ & & \downarrow I \\ \mathcal{O}^G(E, (S_n, d_{S_n})_{n \in \mathbb{N}}) & \xrightarrow{F} & \mathcal{O}^G(E, (X_n, d_{X_n})_{n \in \mathbb{N}}) \\ \downarrow P_k & & \downarrow Q_k \\ \mathcal{O}^G(E, G, d_G) & \xrightarrow{\text{id}} & \mathcal{O}^G(E, G, d_G) \end{array}$$

Explanations follow. The functors \overline{P}_k and \overline{Q}_k are defined as compositions

$$\begin{aligned} \mathcal{O}^G(E, (S_n, d_{S_n})_{n \in \mathbb{N}}) &\rightarrow \mathcal{O}^G(E, S_k, d_{S_k}) \rightarrow \mathcal{O}^G(E, G, d_G) \\ \mathcal{O}^G(E, (X_n, d_{X_n})_{n \in \mathbb{N}}) &\rightarrow \mathcal{O}^G(E, X_k, d_{X_k}) \rightarrow \mathcal{O}^G(E, G, d_G) \end{aligned}$$

where in both cases the first functor is the projection on the k -th factor, and the second functor is induced by the canonical projection $p_k: S_k = G \times \prod_{H \in \mathcal{H}_k} G/\overline{H} \rightarrow G$ and $q_k: X_k = G \times \prod_{H \in \mathcal{H}_k} \text{ind}_{\overline{H}}^G E_H \rightarrow G$ respectively. The functor I is the canonical inclusion. The functor F will be constructed in Proposition 7.1. We have the following facts.

- (i) For all $a \in K_*(\mathcal{O}^G(E, G, d_G))$ and $b \in L_*(\mathcal{O}^G(E, G, d_G))$ there are $\hat{a} \in K_*(\mathcal{O}^G(E, (S_n, d_{S_n})_{n \in \mathbb{N}}))$ and $\hat{b} \in L_*(\mathcal{O}^G(E, (S_n, d_{S_n})_{n \in \mathbb{N}}))$ such that for all k we have $(K_n(P_k))(\hat{a}) = a$ and $(L_*(Q_k))(\hat{b}) = b$. This will be proved in Theorem 6.5.
- (ii) For all k we have $Q_k \circ F = P_k$, see Proposition 7.1.
- (iii) The functor I induces an isomorphism in K - and L -theory. For K -theory this follows from [4, Theorem 7.2]. This result only depends on the properties of K -theory listed in [2, Theorem 5.1]. Since properties are also enjoyed by L -theory, I induces an isomorphism in L -theory as well.

Proof of Theorem 1.2. Because of Theorem 3.1 it suffices to show that the K - and L -theory of $\mathcal{O}^G(E, G, d_G)$ is trivial. Let $a \in K_n(\mathcal{O}^G(E, G, d_G))$ and $b \in L_n(\mathcal{O}^G(E, G, d_G))$. By the first fact there are $\hat{a} \in K_n(\mathcal{O}^G(E, (S_n, d_{S_n})_{n \in \mathbb{N}}))$ and $\hat{b} \in L_n(\mathcal{O}^G(E, (S_n, d_{S_n})_{n \in \mathbb{N}}))$ such that for all k we have $(K_n(P_k))(\hat{a}) = a$ and $(L_n(P_k))(\hat{b}) = b$. It is a consequence of the third fact that for sufficient large k we have $(K_n(Q_k \circ F))(\hat{a}) = 0$ and $(L_n^{-\infty}(Q_k \circ F))(\hat{b}) = 0$. Using the second fact we conclude $a = (K_n(P_k))(\hat{a}) = (K_n(Q_k) \circ F)(\hat{a}) = 0$ and $b = (L_n(P_k))(\hat{b}) = (L_n(Q_k \circ F))(\hat{b}) = 0$. (Compare [4, p.45, Proof of Theorem 1.1].) \square

5. ABSTRACT TRANSFERS FOR ADDITIVE CATEGORIES

Swan group and Dress' equivariant Witt group. We have introduced the additive category with involutions $\mathbf{mod}_{(\mathbb{Z}, G)}$ in Section 2. Recall that it is equivalent to the category of $\mathbb{Z}G$ -modules which are finitely generated free as \mathbb{Z} -modules. We will use the exact structure on $\mathbf{mod}_{(\mathbb{Z}, G)}$ where a sequence is called exact if it is exact as a sequence of $\mathbb{Z}[G]$ -modules (or equivalently as a sequence of abelian groups). Notice that with this exact structure not all exact sequences are split exact over $\mathbb{Z}G$. The *Swan group* and *Dress' equivariant Witt group* are defined with respect to this exact structure as corresponding Grothendieck or Witt groups

$$\text{Sw}(\mathbb{Z}, G) := G_0(\mathbf{mod}_{(\mathbb{Z}, G)}) \quad \text{and} \quad \text{GW}(\mathbb{Z}, G) := W(\mathbf{mod}_{(\mathbb{Z}, G)}),$$

see [18, 6, 14]. Both of these become rings via the tensor product over \mathbb{Z} , equipped with the diagonal G -action, but we will not need this ring structure and ignore it in this paper. By $1_{\text{sw}} \in \text{Sw}(\mathbb{Z}, G)$ we will denote the class of the object given by

the finite set consisting of one element with the trivial G -action and analogously for $1_{\text{GW}} \in \text{GW}(\mathbb{Z}, G)$. In other words, we take the obvious elements whose underlying abelian group is \mathbb{Z} . (These are of course the units for the ring structures.)

For a group homomorphism $\alpha: H \rightarrow G$ there are restriction maps

$$\begin{aligned} \text{res}_\alpha: \text{Sw}(\mathbb{Z}, G) &\rightarrow \text{Sw}(\mathbb{Z}, H) \\ \text{res}_\alpha: \text{GW}(\mathbb{Z}, G) &\rightarrow \text{GW}(\mathbb{Z}, H) \end{aligned}$$

coming from the restriction functor $\text{res}_\alpha: \mathbf{mod}_{(\mathbb{Z}, G)} \rightarrow \mathbf{mod}_{(\mathbb{Z}, H)}$. Clearly, we have $\text{res}_\alpha(1_{\text{Sw}}) = 1_{\text{Sw}}$ and $\text{res}_\alpha(1_{\text{GW}}) = 1_{\text{GW}}$.

For a subgroup $H \subseteq G$ of finite index there are induction homomorphisms

$$\begin{aligned} \text{ind}_H^G: \text{Sw}(\mathbb{Z}, H) &\rightarrow \text{Sw}(\mathbb{Z}, G) \\ \text{ind}_H^G: \text{GW}(\mathbb{Z}, H) &\rightarrow \text{GW}(\mathbb{Z}, G) \end{aligned}$$

coming from the induction functor $\text{ind}_H^G: \mathbf{mod}_{(\mathbb{Z}, H)} \rightarrow \mathbf{mod}_{(\mathbb{Z}, G)}$.

Actually, both $\text{Sw}(\mathbb{Z}, -)$ and $\text{GW}(\mathbb{Z}, -)$ are Green functors. Later on we will make crucial use of the following results due to Swan and Dress.

Theorem 5.1 (Swan [18]; Dress [6]). *Let F be a finite group. Let \mathcal{H} be the family of hyperelementary subgroups of F .*

(i) *There are $\tau_H \in \text{Sw}(\mathbb{Z}, H)$, $H \in \mathcal{H}$ such that*

$$1_{\text{Sw}} = \sum_{H \in \mathcal{H}} \text{ind}_H^F(\tau_H) \in \text{Sw}(\mathbb{Z}, F).$$

(ii) *There are $\sigma_H \in \text{GW}(\mathbb{Z}, H)$, $H \in \mathcal{H}$ such that*

$$1_{\text{GW}} = \sum_{H \in \mathcal{H}} \text{ind}_H^F(\sigma_H) \in \text{GW}(\mathbb{Z}, F).$$

Remark 5.2. In Theorem 5.1 (ii) the family \mathcal{H} can be replaced by the family of subgroups H of F that are either 2-elementary or p -hyperelementary for some prime $p \neq 2$.

Action of $\text{Sw}(\mathbb{Z}, G)$ in K -theory. Let R be a ring and G be a group. Denote by $\mathbf{mod}_{R[G]}$ the category of finitely generated projective $R[G]$ -modules. The tensor product over \mathbb{Z} , equipped with the diagonal G -action, $(T, M) \mapsto T \otimes_{\mathbb{Z}}^{\Delta} M$ defines a bilinear functor

$$-\otimes_{\mathbb{Z}}^{\Delta} -: \mathbf{mod}_{(\mathbb{Z}, G)} \times \mathbf{mod}_{R[G]} \rightarrow \mathbf{mod}_{R[G]}.$$

In particular, we obtain a functor $T \otimes_{\mathbb{Z}}^{\Delta} -: \mathbf{mod}_{R[G]} \rightarrow \mathbf{mod}_{R[G]}$ for every module $T \in \mathbf{mod}_{(\mathbb{Z}, G)}$. Applying K -theory we obtain an endomorphism $K_*(T \otimes_{\mathbb{Z}}^{\Delta} -)$ of $K_*(R[G])$. This endomorphism depends only on the class of T in $\text{Sw}(\mathbb{Z}, G)$ and yields a pairing

$$\mu: \text{Sw}(\mathbb{Z}, G) \otimes K_*(R[G]) \rightarrow K_*(R[G])$$

such that $\mu([T] \otimes a) = K_*(T \otimes_{\mathbb{Z}}^{\Delta} -)(a)$ for all $a \in K_*(R[G])$ (see [16, Corollary 1 on page 106]). This has a generalization as follows. For an additive category \mathcal{B} a functor

$$F: \mathbf{mod}_{(\mathbb{Z}, G)} \times \mathcal{B} \rightarrow \mathcal{B}$$

is said to be *exact* if F is bilinear and for any short exact sequence (which is *not* necessarily split exact) $0 \rightarrow S_0 \xrightarrow{i} S_1 \xrightarrow{p} S_2 \rightarrow 0$ in $\mathbf{mod}_{(\mathbb{Z}, G)}$ and any object B in \mathcal{B} the induced sequence $0 \rightarrow F(S_0, B) \xrightarrow{F(i, \text{id}_B)} F(S_1, B) \xrightarrow{F(p, \text{id}_B)} F(S_2, B) \rightarrow 0$ in $\mathbf{mod}_{(\mathbb{Z}, G)}$ is exact in \mathcal{B} . Recall that a sequence $0 \rightarrow B_0 \xrightarrow{j} B_1 \xrightarrow{q} B_2 \rightarrow 0$ in an additive category \mathcal{B} is called exact if it is split exact, i.e. $q \circ j = 0$ and there exists a morphism $s: B_2 \rightarrow B_0$ such that $q \circ s = \text{id}_{B_2}$ and $j \oplus s: B_0 \oplus B_2 \rightarrow B_1$ is an isomorphism.

Proposition 5.3. *Given an exact functor $F: \mathbf{mod}_{(\mathbb{Z}, G)} \times \mathcal{B} \rightarrow \mathcal{B}$, there is a bilinear pairing*

$$\mu_F: \mathbf{Sw}(\mathbb{Z}, G) \otimes K_*(\mathcal{B}) \rightarrow K_*(\mathcal{B})$$

such that $\mu_F([T] \otimes a) = K_*(F(T, -))(a)$ for all $a \in K_*(\mathcal{B})$.

Action of $\mathbf{GW}(\mathbb{Z}, G)$ in L -theory. Let \mathcal{B} be an additive category with a strict involution $I_{\mathcal{B}}$ and

$$F: \mathbf{mod}_{(R, G)} \times \mathcal{B} \rightarrow \mathcal{B}$$

be an exact functor which is compatible with the involutions, i.e., $I_{\mathcal{B}}(F(-, -)) = F(-^*, I_{\mathcal{B}}(-))$. Then for a module $T \in \mathbf{mod}_{(G, \mathbb{Z})}$ the linear functor $F(T, -): \mathcal{B} \rightarrow \mathcal{B}$ does a priori not induce a map in L -theory because no canonical isomorphism $I_{\mathcal{B}}(F(T, M)) \rightarrow F(T, I_{\mathcal{B}}(M))$ is provided. To fix this, we pick an isomorphism $\varphi: T \rightarrow T^*$ in $\mathbf{mod}_{(\mathbb{Z}, G)}$ such that $\varphi^* = \varphi$, so (T, φ) is a symmetric form in $\mathbf{mod}_{(\mathbb{Z}, G)}$. Then

$$F(\varphi, \mathrm{id}_{I_{\mathcal{B}}(-)}): F(T, I_{\mathcal{B}}(-)) \rightarrow F(T^*, I_{\mathcal{B}}(-)) = I_{\mathcal{B}}(F(T, -))$$

is a natural isomorphism and $F((T, \varphi), -) := (F(T, -), F(\varphi, \mathrm{id}_{I_{\mathcal{B}}(-)})): \mathcal{B} \rightarrow \mathcal{B}$ is a functor of additive categories with involutions. There is the following analog of Proposition 5.3.

Proposition 5.4. *Given an exact functor $F: \mathbf{mod}_{(G, \mathbb{Z})} \times \mathcal{B} \rightarrow \mathcal{B}$ that is compatible with involutions, there is a bilinear pairing*

$$\mu_F: \mathbf{GW}(\mathbb{Z}, G) \otimes L_*(\mathcal{B}) \rightarrow L_*(\mathcal{B})$$

such that $\mu_F([T, \varphi] \otimes b) = L_*(F((T, \varphi), -))(b)$ for all $b \in L_*(\mathcal{B})$ and all symmetric forms (T, φ) over $\mathbf{mod}_{(\mathbb{Z}, G)}$.

Proof. If \mathcal{B} is the category of finitely generated free $R[G]$ -modules and F is the diagonal tensor product, then this is worked out in detail in [6] and [14]. The case of general F and \mathcal{B} is not more complicated. \square

6. THE TRANSFER

Transfer functors. Let G be a group with a metric d_G and E be a G -space. We define a functor

$$\mathrm{tr}: \mathbf{mod}_{(\mathbb{Z}, G)} \times \mathcal{O}^G(E, G, d_G) \rightarrow \mathcal{O}^G(E, G, d_G)$$

as follows. Recall that we have a tensor product functor $\mathbf{mod}_{\mathbb{Z}} \times \mathcal{A} \rightarrow \mathcal{A}$ which is compatible with the involution on $\mathbf{mod}_{\mathbb{Z}}$ and \mathcal{A} , see Section 2. For objects $T = (\mathbb{Z}^n, \rho) \in \mathbf{mod}_{(\mathbb{Z}, G)}$ and $M = (M_z)_{z \in G \times E \times [1, \infty)} \in \mathcal{O}^G(E, G, d_G)$ we define $\mathrm{tr}(T, M) \in \mathcal{O}^G(E, G, d_G)$ by setting

$$(\mathrm{tr}(T, M))_z := \mathbb{Z}^n \otimes_{\mathbb{Z}} M_z$$

for $z \in G \times E \times [1, \infty)$. For morphisms $f \in \mathbf{mod}_{(\mathbb{Z}, G)}$ and $\psi = (\psi_{z, z'})_{z, z' \in G \times E \times [1, \infty)} \in \mathcal{O}^G(E, G, d_G)$ we define $\mathrm{tr}(f, \psi)$ by setting

$$(\mathrm{tr}(f, \psi))_{z, z'} := (f \circ \rho(g^{-1}g')) \otimes_{\mathbb{Z}} \psi_{z, z'}$$

for $z = (g, e, t), z' = (g', e', t') \in G \times E \times [1, \infty)$.

Lemma 6.1. *The functor tr is exact. It is compatible with involutions if \mathcal{A} comes with a (strict) involution.*

Proof. The compatibility with involutions follows from the same compatibility for $\otimes_{\mathbb{Z}}$.

Consider an exact sequence $0 \rightarrow S_0 \xrightarrow{i} S_1 \xrightarrow{q} S_2 \rightarrow 0$ in $\mathbf{mod}_{\mathbb{Z}, G}$. We have to show that for any object M in $\mathcal{O}^G(E, G, d_G)$ that the composite $\mathrm{tr}(q, \mathrm{id}_M) \circ \mathrm{tr}(i, \mathrm{id}_M)$ is trivial, $\mathrm{tr}(q, \mathrm{id}_M): \mathrm{tr}(S_1, M) \rightarrow \mathrm{tr}(S_2, M)$ is split surjective, and that the direct sum of the splitting and the map $\mathrm{tr}(i, \mathrm{id}_M)$ yields an isomorphism $\mathrm{tr}(S_0, M) \oplus \mathrm{tr}(S_2, M) \xrightarrow{\cong} \mathrm{tr}(S_1, M)$. We only construct the splitting of $\mathrm{tr}(q, \mathrm{id}_M)$. Let $s: S \hookrightarrow T$ be a section for q as a map of \mathbb{Z} -modules. Then a section \hat{s} for $\mathrm{tr}(q, \mathrm{id}_M)$ is defined by setting

$$(\hat{s})_{z, z'} := \begin{cases} s \otimes \mathrm{id}_{M_z} & \text{if } z = z' \\ 0 & \text{otherwise.} \end{cases}$$

□

Remark 6.2. To illustrate the proof above consider an epimorphism $p: M \rightarrow N$ of $\mathbb{Z}G$ -modules which are finitely generated free as abelian groups and the induced map of $\mathbb{Z}G$ -modules (with respect to the diagonal action) $p \otimes_{\mathbb{Z}} \mathrm{id}_{\mathbb{Z}G}: M \otimes_{\mathbb{Z}} \mathbb{Z}G \rightarrow N \otimes_{\mathbb{Z}} \mathbb{Z}G$. We want to construct a $\mathbb{Z}G$ -splitting. Choose any map of \mathbb{Z} -modules $s: N \rightarrow M$ with $p \circ s = \mathrm{id}_N$. It exists since we do *not* require that s is compatible with the G -action. Then a $\mathbb{Z}G$ -splitting of $p \otimes_{\mathbb{Z}} \mathrm{id}_{\mathbb{Z}G}$ is given by the $\mathbb{Z}G$ -map $N \otimes_{\mathbb{Z}} \mathbb{Z}G \rightarrow M \otimes_{\mathbb{Z}} \mathbb{Z}G$ sending $n \otimes g$ to $gs(g^{-1}n) \otimes g$.

We will need a variant of tr that combines it with an induction map. This will yield additional control in the target category which is crucial for our argument. Let $\alpha: G \rightarrow F$ be a group homomorphism, H be subgroup of finite index in F . We have defined induction and restriction in Section 2. Consider the functor

$$\mathrm{tr}_\alpha := \mathrm{tr}(\mathrm{res}_\alpha \circ \mathrm{ind}_H^F(-), -): \mathbf{mod}_{(\mathbb{Z}, H)} \times \mathcal{O}^G(E, G, d_G) \rightarrow \mathcal{O}^G(E, G, d_G).$$

Define a quasi-metric $d_{G, H}$ on $G \times G/\overline{H}$ by

$$d_{G, \overline{H}}((g, a\overline{H}), (h, b\overline{H})) := \begin{cases} d_G(g, h) & \text{if } a\overline{H} = b\overline{H}, \\ \infty & \text{otherwise.} \end{cases}$$

The projection $p_H: G \times G/\overline{H} \rightarrow G$ induces a functor $P_H: \mathcal{O}^G(E, G \times G/\overline{H}, d_{G, \overline{H}}) \rightarrow \mathcal{O}^G(E, G, d_G)$ and we will see that we can lift tr_α against P_H . Define a functor

$$\tilde{\mathrm{tr}}_\alpha: \mathbf{mod}_{(\mathbb{Z}, H)} \times \mathcal{O}^G(E, G, d_G) \rightarrow \mathcal{O}^G(E, G \times G/\overline{H}, d_{G, \overline{H}})$$

as follows. For objects $T = (\mathbb{Z}^n, \rho) \in \mathbf{mod}_{(\mathbb{Z}, H)}$ and $M = (M_z)_{z \in G \times E \times [1, \infty)} \in \mathcal{O}^G(E, G, d_G)$ we define $\tilde{\mathrm{tr}}_\alpha(T, M)$ by setting

$$(\tilde{\mathrm{tr}}_\alpha(T, M))_y := \mathbb{Z}^n \otimes_{\mathbb{Z}} M_z$$

for $y = (g, a\overline{H}, e, t) \in G \times G/\overline{H}, E \times [1, \infty)$ and $z := (g, e, t)$. For morphisms $f \in \mathbf{mod}_{(\mathbb{Z}, H)}$ and $\psi = (\psi_{z, z'})_{z, z' \in G \times E \times [1, \infty)} \in \mathcal{O}^G(E, G, d_G)$ we define $\tilde{\mathrm{tr}}_\alpha(f, \psi)$ by setting

$$(\tilde{\mathrm{tr}}_\alpha(f, \psi))_{y, y'} := \begin{cases} f \circ \rho(\alpha(g^{-1}g')) \otimes_{\mathbb{Z}} \psi_{z, z'} & \text{if } \alpha(g^{-1}g') \in H, \\ 0 & \text{otherwise.} \end{cases}$$

for $y = (g, a\overline{H}, e, t), y' = (g', a'\overline{H}, e', t') \in G \times G/\overline{H}, E \times [1, \infty)$ and $z := (g, e, t), z' := (g', e', t')$. (The extra G/\overline{H} -factor incorporates the induction from H to F ; the appearance of α incorporates the restriction along α .)

The following Lemma is a simple exercise in the definitions of tr_α and $\tilde{\mathrm{tr}}_\alpha$.

Lemma 6.3.

(i) $P_H \circ \tilde{\mathrm{tr}}_\alpha$ and tr_α are equivalent functors.

(ii) if ψ is an R -controlled morphism in $\mathcal{O}^G(E, G, d_G)$ and $f \in \text{mod}_{(\mathbb{Z}, H)}$ is any morphism, then $\tilde{\text{tr}}_\alpha(f, \psi)$ is R -controlled in $\mathcal{O}^G(E, G, d_G)$.

Surjectivity of the P_k in (4.1). In the remainder of this section we use the notation from Section 4. In particular G will from now on be a Farrell-Hsiang group. We denote by $(p_n)_*: \mathcal{O}^G(E, S_n, d_{S_n}) \rightarrow \mathcal{O}^G(E, G, d_G)$ the functor induced by the projection $p_n: S_n = G \times \coprod_{H \in \mathcal{H}_n} G/\overline{H} \rightarrow G$.

Proposition 6.4. *Let $n \in \mathbb{N}$.*

- (i) *There are linear functors $F_n^+, F_n^-: \mathcal{O}^G(E, G, d_G) \rightarrow \mathcal{O}^G(E, S_n, d_{S_n})$ with the following two properties*
- $K_*((p_n)_* \circ F_n^+) - K_*((p_n)_* \circ F_n^-)$ *is the identity on $K_*(\mathcal{O}^G(E, G, d_G))$;*
 - *if $R > 0$ and $\psi \in \mathcal{O}^G(E, G, d_G)$ is R -controlled, then $F_n^+(\psi)$ and $F_n^-(\psi)$ are both also R -controlled.*
- (ii) *There are functors of additive categories with involutions $G_n^+ = (G_n^+, E_n^+)$ and $G_n^- = (G_n^-, E_n^-): \mathcal{O}^G(E, G, d_G) \rightarrow \mathcal{O}^G(E, S_n, d_{S_n})$ with the following properties*
- $L_*((p_n)_* \circ G_n^+) - L_*((p_n)_* \circ G_n^-)$ *is the identity on $L_*(\mathcal{O}^G(E, G, d_G))$;*
 - *if $R > 0$ and $\psi \in \mathcal{O}^G(E, G, d_G)$ is R -controlled, then $G_n^+(\psi)$ and $G_n^-(\psi)$ are both also R -controlled.*
 - *Denote by I both the involution on $\mathcal{O}^G(E, S_n, d_{S_n})$ and the involution on $\mathcal{O}^G(E, G, d_G)$. For each object $M \in \mathcal{O}^G(E, G, d_G)$, the isomorphisms $E_n^+(M): G_n^+(I(M)) \rightarrow I(G_n^+(M))$ and $E_n^-(M): G_n^-(I(M)) \rightarrow I(G_n^-(M))$ are 0-controlled.*

Proof. (i) By Theorem 5.1 (i) there are $\tau_H \in \text{Sw}(\mathbb{Z}, H)$, $H \in \mathcal{H}_n$ such that $1_{\text{Sw}} = \sum_{H \in \mathcal{H}_n} \text{ind}_H^{F_n}(\tau_H) \in \text{Sw}(\mathbb{Z}, F_n)$. Any element in $\text{Sw}(\mathbb{Z}, H)$ can be written as the difference of the classes of two modules. Pick modules T_H^+ and $T_H^- \in \text{mod}_{(\mathbb{Z}, G)}$, $H \in \mathcal{H}_n$ such that $\tau_H = [T_H^+] - [T_H^-]$. Because res_{α_n} sends $1_{\text{Sw}} \in \text{Sw}(\mathbb{Z}, F_n)$ to $1_{\text{Sw}} \in \text{Sw}(\mathbb{Z}, G)$ we obtain

$$1_{\text{Sw}} = \sum_{H \in \mathcal{H}_n} [\text{res}_{\alpha_n} \circ \text{ind}_H^{F_n}(T_H^+)] - [\text{res}_{\alpha_n} \circ \text{ind}_H^{F_n}(T_H^-)] \in \text{Sw}(\mathbb{Z}, G)$$

For $H \in \mathcal{H}_n$ we have a canonical inclusion $G \times G/\overline{H} \rightarrow S_n = G \times \coprod_{K \in \mathcal{H}_n} G/\overline{K}$ that induces an inclusion $\mathcal{O}^G(E, G \times G/\overline{H}, d_{G, \overline{H}}) \rightarrow \mathcal{O}^G(E, S_n, d_{S_n})$. Define F_H^\pm as the composition of $\tilde{\text{tr}}_{\alpha_n}(T_H^\pm, -)$ with this inclusion. Then $K_*((p_n)_* \circ F^\pm H) = K_*(\text{tr}(\text{res}_{\alpha_n} \text{ind}_H^{F_n}(T_H^\pm), -))$ by Lemma 6.3 (i). Define now

$$F_n^\pm := \bigoplus_{H \in \mathcal{H}_n} F_H^\pm.$$

The functor tr is exact by Lemma 6.1 and so Proposition 5.3 applies. Therefore we can compute for all $a \in K_*(\mathcal{O}^G(E, G, d_G))$

$$\begin{aligned} & K_*((p_n)_* \circ F_n^+)(a) - K_*((p_n)_* \circ F_n^-)(a) \\ &= \sum_{H \in \mathcal{H}_n} K_*((p_n)_* \circ F_H^+)(a) - K_*((p_n)_* \circ F_H^-)(a) \\ &= \sum_{H \in \mathcal{H}_n} K_*(\text{tr}(\text{res}_{\alpha_n} \circ \text{ind}_H^{F_n}(T_H^+)))(a) - K_*(\text{tr}(\text{res}_{\alpha_n} \circ \text{ind}_H^{F_n}(T_H^-)))(a) \\ &= \sum_{H \in \mathcal{H}_n} \mu_{\text{tr}}([\text{res}_{\alpha_n} \circ \text{ind}_H^{F_n}(T_H^+)] \otimes a) - \mu_{\text{tr}}([\text{res}_{\alpha_n} \circ \text{ind}_H^{F_n}(T_H^-)] \otimes a) \\ &= \mu_{\text{tr}}((\sum_{H \in \mathcal{H}_n} [\text{res}_{\alpha_n} \circ \text{ind}_H^{F_n}(T_H^+)] - [\text{res}_{\alpha_n} \circ \text{ind}_H^{F_n}(T_H^-)]) \otimes a) \\ &= \mu_{\text{tr}}(1_{\text{Sw}} \otimes a) = a \end{aligned}$$

If $R > 0$ and $\psi \in \mathcal{O}^G(E, G, d_G)$ is R -controlled then each $F_H^\pm(\psi)$ is R -controlled, because of the control property of $\tilde{\text{tr}}_{\alpha_n}$ (Lemma 6.3 (ii)) and because $G \times G/\overline{H} \rightarrow S_n$ is an isometric embedding. The direct sum of R -controlled morphisms is again R -controlled and therefore $F^+(\psi)$ and $F^-(\psi)$ are both R -controlled.

(ii) We can proceed exactly as in the K -theory case. By Theorem 5.1 (ii) there are $\sigma_H \in \text{GW}(\mathbb{Z}, H)$, $H \in \mathcal{H}_n$ such that $1_{\text{GW}} = \sum_{H \in \mathcal{H}_n} \text{ind}_{F_n}^H(\sigma_H)$. Any element in $\text{GW}(\mathbb{Z}, H)$ can be written as the difference of the classes of two symmetric forms. Pick symmetric forms (T_H^+, φ_H^+) and (T_H^-, φ_H^-) over $\text{mod}_{(\mathbb{Z}, G)}$, $H \in \mathcal{H}_n$ such that $\sigma_H = [(T_H^+, \varphi_H^+) - (T_H^-, \varphi_H^-)]$. Define G_H^\pm as the composition of $\text{tr}_{\alpha_n}((T_H^\pm, \varphi_H^\pm), -)$ with the inclusion $\iota_H: \mathcal{O}^G(E, G \times G/\overline{H}, d_{G, \overline{H}}) \rightarrow \mathcal{O}^G(E, S_n, d_{S_n})$ and set

$$G_n^\pm := \bigoplus_{H \in \mathcal{H}_n} G_H^\pm.$$

As in the K -theory case it follows (using now Proposition 5.4) that for all $b \in L_*(\mathcal{O}^G(E, G, d_G))$ we have

$$L_*((p_n)_* \circ G_n^+)(b) - L_*((p_n)_* \circ G_n^-)(b) = b$$

and that $G_n^\pm(\psi)$ is R -controlled, whenever ψ is R -controlled.

It remains to prove the final claim. Let M be an object from $\mathcal{O}^G(E, G, d_G)$. Then

$$\begin{aligned} G_n^\pm(I(M)) &= \bigoplus_H G_H^\pm(I(M)) = \bigoplus_H \iota_H(\tilde{\text{tr}}_{\alpha_n}(T_H^\pm, I(M))) \\ I(G_n^\pm(M)) &= \bigoplus_H I(G_H^\pm(M)) = \bigoplus_H I(\iota_H(\tilde{\text{tr}}_{\alpha_n}(T_H^\pm, M))) \\ &= \bigoplus_H \iota_H(\tilde{\text{tr}}_{\alpha_n}((T_H^\pm)^*, I(M))) \end{aligned}$$

and

$$E_n^\pm(M) = \bigoplus_H \iota_H(\tilde{\text{tr}}_{\alpha_n}(\varphi_H^\pm, \text{id}_{I(M)})).$$

The control claim follows from Lemma 6.3 (ii) because $\text{id}_{I(M)}$ is 0-controlled. \square

Theorem 6.5.

(i) For all $a \in K_*(\mathcal{O}^G(E, G, d_G))$ there is $\hat{a} \in K_*(\mathcal{O}^G(E, (S_n, d_{S_n})_{n \in \mathbb{N}}))$ such that for all k we have $(K_*(P_k))(\hat{a}) = a$.

(ii) For all $b \in L_*(\mathcal{O}^G(E, G, d_G))$ there is $\hat{b} \in L_*(\mathcal{O}^G(E, (S_n, d_{S_n})_{n \in \mathbb{N}}))$ such that for all k we have $(L_*(P_k))(\hat{b}) = b$.

Proof. (i) Let F_n^+, F_n^- be the sequences of functors from Proposition 6.4 (i). Because of the control property in 6.4 (i) the product functors

$$\prod_n F_n^+ : \mathcal{O}^G(E, G, d_G) \rightarrow \prod_n \mathcal{O}^G(E, S_n, d_{S_n})$$

lift uniquely to functors

$$F^\pm : \mathcal{O}^G(E, G, d_G) \rightarrow \mathcal{O}^G(E, (S_n, d_{S_n})_{n \in \mathbb{N}}).$$

Then $P_k \circ F^\pm = (p_k)_* \circ F_k^\pm$ for all $k \in \mathbb{N}$. Thus the first assertion in 6.4 (i) implies that $K_*(P_k)(K_*(F^+)(a) - K_*(F^-)(a)) = a$ for all $a \in \mathcal{O}^G(E, G, d_G)$. Therefore we can set $\hat{a} := K_*(F^+)(a) - K_*(F^-)(a)$.

(ii) For L -theory we can argue exactly as we did for K -theory, now using the G_n^\pm from Proposition 6.4 (ii). Here the third assertion in 6.4 (ii) is needed to ensure that the E_n^\pm can be combined to a natural transformation, just as the second assertion is needed to ensure that the G_n^\pm can be combined to a functor. \square

7. THE FUNCTOR F

We use the notation from Section 4. Note first that for any subgroup U of G there is a bijection of G -sets $G \times G/U \rightarrow \text{ind}_G^U \text{res}_U^G G = G \times_U G$ defined by $(a, gU) \mapsto (g, g^{-1}a)$; the inverse is given by $(g, b) \mapsto gb, gU$. (We use the diagonal G -action on $G \times G/U$.) For $H \in \mathcal{H}_n$ we obtain a G -map $\tilde{f}_H: G \times G/\overline{H} \rightarrow \text{ind}_G^{\overline{H}} E_H$ by composing this bijection (for $U = \overline{H}$) with $\text{ind}_G^{\overline{H}} f_H: \text{ind}_G^{\overline{H}} G \rightarrow \text{ind}_G^{\overline{H}} E_H$. Define the G -map $f_n: S_n \rightarrow X_n$ by

$$f_n(a, g\overline{H}) := (a, \tilde{f}_H(a, g\overline{H}) = (a, g, f_H(g^{-1}a)),$$

for $a, g \in G$ and $H \in \mathcal{H}_n$.

Proposition 7.1. *The sequence of maps $(f_n)_{n \in \mathbb{N}}$ induces a functor*

$$F: \mathcal{O}^G(E, (S_n, d_{S_n})_{n \in \mathbb{N}}) \rightarrow \mathcal{O}^G(E, (X_n, d_{X_n})_{n \in \mathbb{N}}).$$

For all k we have $q_k \circ F = p_k$.

Proof. We need to show that the sequence $(f_n)_{n \in \mathbb{N}}$ is compatible with the metric control conditions for the sequences of quasi-metrics $(d_{S_n})_{n \in \mathbb{N}}$ and $(d_{X_n})_{n \in \mathbb{N}}$ more precisely we need to show that for any $r \in (0, \infty)$ there is $R \in (0, \infty)$ such that for all n and $s, s' \in S_n$ the implication

$$(7.2) \quad d_{S_n}(s, s') < r \quad \implies \quad d_{X_n}(f_n(s), f_n(s')) < R$$

holds.

Let $r \in (0, \infty)$ be given. The G -action on S_n is cofinite, the quasi-metrics d_{S_n} and d_{G_n} are G -invariant and f_n is G -equivariant. For each $s \in S_n$ there are only finitely many $s' \in S_n$ such that $d_{S_n}(s', s) < r$, because the word metric d_G has this property on G . This implies that $D_r := \{d_{X_n}(f_n(s), f_n(s')) \mid n < r, s, s' \in S_n, d_{S_n}(s, s') < r\}$ is a finite set. We can therefore define $R := 1 + r + \max D_r$. We claim that then (7.2) holds for all n and all $s, s' \in S_n$. If $n < r$, then this is clear from the definition of R . Let $n > r$ and $s, s' \in S_n$ with $d_{S_n}(s, s') < r$. Write $s = (a, g\overline{H})$ and $s' = (a', g'\overline{H}')$ with $H, H' \in \mathcal{H}_n$, $a, a', g, g' \in G$. Since $d_{S_n}(s, s') < r < \infty$ it follows from the definition of d_{S_n} that $H = H'$, $g\overline{H} = g'\overline{H}'$ and $d_G(a, a') < r < n$. Since d_G is G -invariant we also have $d_G(g^{-1}a, g^{-1}a') < n$. We conclude from the crucial contracting property of f_H that $d_{E_H}^1(f_H(g^{-1}a), f_H(g^{-1}a')) < \frac{1}{n}$. Since $s = (a, g\overline{H})$, $s' = (a', g'\overline{H}')$ we have $f_n(s) = (a, g, f_H(g^{-1}a))$, $f_n(s') = (a', g, f_H(g^{-1}a'))$. Thus

$$\begin{aligned} d_{X_n}(f_n(s), f_n(s')) &= d_{X_n}((a, g, f_H(g^{-1}a)), (a', g, f_H(g^{-1}a'))) \\ &= d_G(a, a') + n \cdot d_{\text{ind}_G^{\overline{H}} E_H}^1((g, f_H(g^{-1}a)), (g, f_H(g^{-1}a'))) \\ &= d_G(a, a') + n \cdot d_{E_H}^1(f_H(g^{-1}a), f_H(g^{-1}a')) \\ &< r + n \cdot \frac{1}{n} = r + 1 < R. \end{aligned}$$

This proves our claim. Thus $(f_n)_{n \in \mathbb{N}}$ induces a functor F .

For the canonical projections $p_k: S_k \rightarrow G$ and $q_k: X_k \rightarrow G$ we have $q_k \circ f_k = p_k$. This implies that $Q_k \circ F = P_k$. \square

REFERENCES

- [1] A. Bartels, F. T. Farrell and W. Lück. The Farrell-Jones Conjecture for cocompact lattices in virtually connected Lie groups. preprint, 2011.
- [2] A. Bartels and W. Lück. The Borel conjecture for hyperbolic and CAT(0)-groups. Preprintreihe SFB 478 — Geometrische Strukturen in der Mathematik, Heft 506 Münster, arXiv:0901.0442v1 [math.GT], 2009.
- [3] A. Bartels and W. Lück. On twisted group rings with twisted involutions, their module categories and L -theory. In *Cohomology of groups and algebraic K-theory*, volume 12 of *Advanced Lectures in Mathematics*, pages 1–55, Somerville, U.S.A., 2009. International Press.

- [4] A. Bartels, W. Lück, and H. Reich. The K -theoretic Farrell-Jones conjecture for hyperbolic groups. *Invent. Math.*, 172(1):29–70, 2008.
- [5] A. Bartels, W. Lück, and H. Reich. On the Farrell-Jones Conjecture and its applications. *Journal of Topology*, 1:57–86, 2008.
- [6] A. W. M. Dress. Induction and structure theorems for orthogonal representations of finite groups. *Ann. of Math. (2)*, 102(2):291–325, 1975.
- [7] F. T. Farrell and W. C. Hsiang. The topological-Euclidean space form problem. *Invent. Math.*, 45(2):181–192, 1978.
- [8] F. T. Farrell and W. C. Hsiang. The Whitehead group of poly-(finite or cyclic) groups. *J. London Math. Soc. (2)*, 24(2):308–324, 1981.
- [9] F. T. Farrell and W. C. Hsiang. Topological characterization of flat and almost flat Riemannian manifolds M^n ($n \neq 3, 4$). *Amer. J. Math.*, 105(3):641–672, 1983.
- [10] F. T. Farrell and L. E. Jones. K -theory and dynamics. II. *Ann. of Math. (2)*, 126(3):451–493, 1987.
- [11] F. T. Farrell and L. E. Jones. The surgery L -groups of poly-(finite or cyclic) groups. *Invent. Math.*, 91(3):559–586, 1988.
- [12] F. T. Farrell and L. E. Jones. Isomorphism conjectures in algebraic K -theory. *J. Amer. Math. Soc.*, 6(2):249–297, 1993.
- [13] F. T. Farrell and L. E. Jones. Topological rigidity for compact non-positively curved manifolds. In *Differential geometry: Riemannian geometry (Los Angeles, CA, 1990)*, pages 229–274. Amer. Math. Soc., Providence, RI, 1993.
- [14] W. Lück and A. A. Ranicki. Surgery transfer. In *Algebraic topology and transformation groups (Göttingen, 1987)*, pages 167–246. Springer-Verlag, Berlin, 1988.
- [15] W. Lück and H. Reich. The Baum-Connes and the Farrell-Jones conjectures in K - and L -theory. In *Handbook of K -theory. Vol. 1, 2*, pages 703–842. Springer, Berlin, 2005.
- [16] D. Quillen. Higher algebraic K -theory. I. In *Algebraic K -theory, I: Higher K -theories (Proc. Conf., Battelle Memorial Inst., Seattle, Wash., 1972)*, pages 85–147. Lecture Notes in Math., Vol. 341. Springer-Verlag, Berlin, 1973.
- [17] F. Quinn. Hyperelementary assembly for K -theory of virtually abelian groups. Preprint, arXiv:math.KT/0509294, 2005.
- [18] R. G. Swan. Induced representations and projective modules. *Ann. of Math. (2)*, 71:552–578, 1960.

WESTFÄLISCHE WILHELMS-UNIVERSITÄT MÜNSTER, MATHEMATISCHES INSTITUT, EINSTEINSTR. 62,
D-48149 MÜNSTER, GERMANY

E-mail address: bartelsa@math.uni-muenster.de
URL: <http://www.math.uni-muenster.de/u/bartelsa>

MATHEMATISCHES INSTITUT DER UNIVERSITÄT BONN, ENDENICHER ALLEE 60, 53115 BONN,
GERMANY

E-mail address: wolfgang.lueck@him.uni-bonn.de
URL: <http://www.him.uni-bonn.de/lueck>