

The β -function in duality-covariant noncommutative ϕ^4 -theory

Harald GROSSE¹ and Raimar WULKENHAAR²

¹ *Institut für Theoretische Physik, Universität Wien
Boltzmannngasse 5, A-1090 Wien, Austria*

² *Max-Planck-Institut für Mathematik in den Naturwissenschaften
Inselstraße 22-26, D-04103 Leipzig, Germany*

Abstract

We compute the one-loop β -functions describing the renormalisation of the coupling constant λ and the frequency parameter Ω for the real four-dimensional duality-covariant noncommutative ϕ^4 -model, which is renormalisable to all orders. The contribution from the one-loop four-point function is reduced by the one-loop wavefunction renormalisation, but the β_λ -function remains non-negative. Both β_λ and β_Ω vanish at the one-loop level for the duality-invariant model characterised by $\Omega = 1$. Moreover, β_Ω also vanishes in the limit $\Omega \rightarrow 0$, which defines the standard noncommutative ϕ^4 -quantum field theory. Thus, the limit $\Omega \rightarrow 0$ exists at least at the one-loop level.

¹harald.grosse@univie.ac.at

²raimar.wulkenhaar@mis.mpg.de

1 Introduction

For many years, the renormalisation of quantum field theories on noncommutative \mathbb{R}^4 has been an open problem [1]. Recently, we have proven in [2] that the real duality-covariant ϕ^4 -model on noncommutative \mathbb{R}^4 is renormalisable to all orders. The duality transformation exchanges positions and momenta [3],

$$\hat{\phi}(p) \leftrightarrow \pi^2 \sqrt{|\det \theta|} \phi(x), \quad p_\mu \leftrightarrow \tilde{x}_\mu := 2(\theta^{-1})_{\mu\nu} x^\nu, \quad (1)$$

where $\hat{\phi}(p_a) = \int d^4x e^{(-1)^a i p_a \cdot \mu x_a} \phi(x_a)$. The subscript a refers to the cyclic order in the \star -product. The duality-covariant noncommutative ϕ^4 -action is given by

$$S[\phi; \mu_0, \lambda, \Omega] := \int d^4x \left(\frac{1}{2} (\partial_\mu \phi) \star (\partial^\mu \phi) + \frac{\Omega^2}{2} (\tilde{x}_\mu \phi) \star (\tilde{x}^\mu \phi) + \frac{\mu_0^2}{2} \phi \star \phi + \frac{\lambda}{4!} \phi \star \phi \star \phi \star \phi \right) (x). \quad (2)$$

Under the transformation (1) one has

$$S[\phi; \mu_0, \lambda, \Omega] \mapsto \Omega^2 S\left[\phi; \frac{\mu_0}{\Omega}, \frac{\lambda}{\Omega^2}, \frac{1}{\Omega}\right]. \quad (3)$$

In the special case $\Omega = 1$ the action $S[\phi; \mu_0, \lambda, 1]$ is invariant under the duality (1). Moreover, $S[\phi; \mu_0, \lambda, 1]$ can be written as a standard matrix model which is closely related to an exactly solvable model [4].

Knowing that the action (2) gives rise to a renormalisable quantum field theory [2], it is interesting to compute the β_λ and β_Ω functions which describe the renormalisation of the coupling constant λ and of the oscillator frequency Ω . Whereas we have proven the renormalisability in the Wilson-Polchinski approach [5, 6] adapted to non-local matrix models [7], we compute the one-loop β_λ and β_Ω functions by standard Feynman graph calculations. Of course, these are Feynman graphs parametrised by matrix indices instead of momenta. We rely heavily on the power-counting behaviour proven in [2], which allows us to ignore in the β -functions all non-planar graphs and the detailed index dependence of the planar two- and four-point graphs. Thus, only the lowest-order (discrete) Taylor expansion of the planar two- and four-point graphs can contribute to the β -functions. This means that we cannot refer to the usual symmetry factors of commutative ϕ^4 -theory so that we have to carefully recompute the graphs.

We obtain interesting consequences for the limiting cases $\Omega = 1$ and $\Omega = 0$ as discussed in Section 5.

2 Definition of the model

The noncommutative \mathbb{R}^4 is defined as the algebra \mathbb{R}_θ^4 which as a vector space is given by the space $\mathcal{S}(\mathbb{R}^4)$ of (complex-valued) Schwartz class functions of rapid decay, equipped

with the multiplication rule

$$(a \star b)(x) = \int \frac{d^4 k}{(2\pi)^4} \int d^4 y a(x + \frac{1}{2}\theta \cdot k) b(x+y) e^{ik \cdot y} , \quad (4)$$

$$(\theta \cdot k)^\mu = \theta^{\mu\nu} k_\nu , \quad k \cdot y = k_\mu y^\mu , \quad \theta^{\mu\nu} = -\theta^{\nu\mu} .$$

We place ourselves into a coordinate system in which the only non-vanishing components $\theta_{\mu\nu}$ are $\theta_{12} = -\theta_{21} = \theta_{34} = -\theta_{42} = \theta$. We use an adapted base

$$b_{mn}(x) = f_{m^1 n^1}(x^1, x^2) f_{m^2 n^2}(x^3, x^4) , \quad m = \frac{m^1}{m^2} \in \mathbb{N}^2 , \quad n = \frac{n^1}{n^2} \in \mathbb{N}^2 , \quad (5)$$

where the base $f_{m^1 n^1}(x^1, x^2) \in \mathbb{R}_\theta^2$ is given in [8]. This base satisfies

$$(b_{mn} \star b_{kl})(x) = \delta_{nk} b_{ml}(x) , \quad \int d^4 x b_{mn}(x) = 4\pi^2 \theta^2 \delta_{mn} . \quad (6)$$

According to [2], the duality-covariant ϕ^4 -action (2) expands as follows in the matrix base (5):

$$S[\phi; \mu_0, \lambda, \Omega] = 4\pi^2 \theta^2 \sum_{m,n,k,l \in \mathbb{N}^2} \left(\frac{1}{2} G_{mn;kl} \phi_{mn} \phi_{kl} + \frac{\lambda}{4!} \phi_{mn} \phi_{nk} \phi_{kl} \phi_{lm} \right) , \quad (7)$$

where $\phi(x) = \sum_{m,n \in \mathbb{N}^2} \phi_{mn} b_{mn}(x)$ and

$$G_{mn;kl} = \left(\mu_0^2 + \frac{2}{\theta} (1 + \Omega^2) (m^1 + n^1 + m^2 + n^2 + 2) \right) \delta_{n^1 k^1} \delta_{m^1 l^1} \delta_{n^2 k^2} \delta_{m^2 l^2}$$

$$- \frac{2}{\theta} (1 - \Omega^2) \left((\sqrt{(n^1+1)(m^1+1)}) \delta_{n^1+1, k^1} \delta_{m^1+1, l^1} + \sqrt{n^1 m^1} \delta_{n^1-1, k^1} \delta_{m^1-1, l^1} \right) \delta_{n^2 k^2} \delta_{m^2 l^2}$$

$$+ \left(\sqrt{(n^2+1)(m^2+1)} \delta_{n^2+1, k^2} \delta_{m^2+1, l^2} + \sqrt{n^2 m^2} \delta_{n^2-1, k^2} \delta_{m^2-1, l^2} \right) \delta_{n^1 k^1} \delta_{m^1 l^1} . \quad (8)$$

The quantum field theory is defined by the partition function

$$Z[J] = \int \left(\prod_{a,b \in \mathbb{N}^2} d\phi_{ab} \right) \exp \left(- S[\phi] - 4\pi^2 \theta^2 \sum_{m,n \in \mathbb{N}^2} \phi_{mn} J_{nm} \right) . \quad (9)$$

For the free theory defined by $\lambda = 0$ in (7), the solution of (9) is given by

$$Z[J]|_{\lambda=0} = Z[0] \exp \left(4\pi^2 \theta^2 \sum_{m,n,k,l \in \mathbb{N}^2} \frac{1}{2} J_{mn} \Delta_{mn;kl} J_{kl} \right) , \quad (10)$$

where the propagator Δ is defined as the inverse of the kinetic matrix G :

$$\sum_{k,l \in \mathbb{N}^2} G_{mn;kl} \Delta_{lk;sr} = \sum_{\in \mathbb{N}^2} \Delta_{nm;lk} G_{kl;rs} = \delta_{mr} \delta_{ns} . \quad (11)$$

We have derived the propagator in [2]:

$$\begin{aligned}
\Delta_{\substack{m^1 & n^1 & k^1 & l^1 \\ m^2 & n^2 & k^2 & l^2}} &= \frac{\theta}{2(1+\Omega)^2} \delta_{m^1+k^1, n^1+l^1} \delta_{m^2+k^2, n^2+l^2} \\
&\times \sum_{v^1=\frac{|m^1-l^1|}{2}}^{\frac{\min(m^1+l^1, n^1+k^1)}{2}} \sum_{v^2=\frac{|m^2-l^2|}{2}}^{\frac{\min(m^2+l^2, n^2+k^2)}{2}} B\left(1+\frac{\mu_0^2\theta}{8\Omega}+\frac{1}{2}(m^1+m^2+k^1+k^2)-v^1-v^2, 1+2v^1+2v^2\right) \\
&\times {}_2F_1\left(\begin{matrix} 1+2v^1+2v^2, \frac{\mu_0^2\theta}{8\Omega}-\frac{1}{2}(m^1+m^2+k^1+k^2)+v^1+v^2 \\ 2+\frac{\mu_0^2\theta}{8\Omega}+\frac{1}{2}(m^1+m^2+k^1+k^2)+v^1+v^2 \end{matrix} \middle| \frac{(1-\Omega)^2}{(1+\Omega)^2}\right) \\
&\times \prod_{i=1}^2 \sqrt{\binom{n^i}{v^i+\frac{n^i-k^i}{2}} \binom{k^i}{v^i+\frac{k^i-n^i}{2}} \binom{m^i}{v^i+\frac{m^i-l^i}{2}} \binom{l^i}{v^i+\frac{l^i-m^i}{2}}} \left(\frac{(1-\Omega)^2}{(1+\Omega)^2}\right)^{v^i}}. \tag{12}
\end{aligned}$$

Here, $B(a, b)$ is the Beta-function and ${}_2F_1\left(\begin{smallmatrix} a, b \\ c \end{smallmatrix} \middle| z\right)$ the hypergeometric function.

As usual we solve the interacting theory perturbatively:

$$\begin{aligned}
Z[J] &= Z[0] \exp\left(-V\left[\frac{\partial}{\partial J}\right]\right) \exp\left(4\pi^2\theta^2 \sum_{m,n,k,l \in \mathbb{N}^2} \frac{1}{2} J_{mn} \Delta_{mn;kl} J_{kl}\right), \\
V\left[\frac{\partial}{\partial J}\right] &:= \frac{\lambda}{4!(4\pi^2\theta^2)^3} \sum_{m,n,k,l \in \mathbb{N}^2} \frac{\partial^4}{\partial J_{ml} \partial J_{lk} \partial J_{kn} \partial J_{nm}}. \tag{13}
\end{aligned}$$

It is convenient to pass to the generating functional of connected Green's functions, $W[J] = \ln Z[J]$:

$$\begin{aligned}
W[J] &= \ln Z[0] + W_{\text{free}}[J] + \ln\left(1 + e^{-W_{\text{free}}[J]} \left(\exp\left(-V\left[\frac{\partial}{\partial J}\right]\right) - 1\right) e^{W_{\text{free}}[J]}\right), \\
W_{\text{free}}[J] &:= 4\pi^2\theta^2 \sum_{m,n,k,l \in \mathbb{N}^2} \frac{1}{2} J_{mn} \Delta_{mn;kl} J_{kl}. \tag{14}
\end{aligned}$$

In order to obtain the expansion in λ one has to expand $\ln(1+x)$ as a power series in x and $\exp(-V)$ as a power series in V . By Legendre transformation we pass to the generating functional of one-particle irreducible (1PI) Green's functions:

$$\Gamma[\phi^{cl}] := 4\pi^2\theta^2 \sum_{m,n \in \mathbb{N}^2} \phi_{mn}^{cl} J_{nm} - W[J], \tag{15}$$

where J has to be replaced by the inverse solution of

$$\phi_{mn}^{cl} := \frac{1}{4\pi^2\theta^2} \frac{\partial W[J]}{\partial J_{nm}}. \tag{16}$$

3 Renormalisation group equation

The computation of the expansion coefficients

$$\Gamma_{m_1 n_1; \dots; m_N n_N} := \frac{1}{N!} \frac{\partial^N \Gamma[\phi^{cl}]}{\partial \phi_{m_1 n_1}^{cl} \dots \partial \phi_{m_N n_N}^{cl}} \quad (17)$$

of the effective action involves possibly divergent sums over undetermined loop indices. Therefore, we have to introduce a cut-off \mathcal{N} for all loop indices. According to [2], the expansion coefficients (17) can be decomposed into a relevant/marginal and an irrelevant piece. As a result of the renormalisation proof, the relevant/marginal parts have—after a rescaling of the field amplitude—the same form as the initial action (2), (7) and (8), now parametrised by the “physical” mass, coupling constant and oscillator frequency:

$$\Gamma_{\text{rel/marg}}[\mathcal{Z}\phi^{cl}] = S[\phi^{cl}; \mu_{\text{phys}}, \lambda_{\text{phys}}, \Omega_{\text{phys}}] . \quad (18)$$

In the renormalisation process, the physical quantities μ_{phys}^2 , λ_{phys} and Ω_{phys} are kept constant with respect to the cut-off \mathcal{N} . This is achieved by starting from a carefully adjusted initial action $S[\mathcal{Z}[\mathcal{N}]\phi, \mu_0[\mathcal{N}], \lambda[\mathcal{N}], \Omega[\mathcal{N}]]$, which gives rise to the bare effective action $\Gamma[\phi^{cl}; \mu_0[\mathcal{N}], \lambda[\mathcal{N}], \Omega[\mathcal{N}], \mathcal{N}]$. Expressing the bare parameters μ_0, λ, Ω as a function of the physical quantities and the cut-off, the expansion coefficients of the renormalised effective action

$$\Gamma^R[\phi^{cl}; \mu_{\text{phys}}, \lambda_{\text{phys}}, \Omega_{\text{phys}}] := \Gamma[\mathcal{Z}[\mathcal{N}]\phi^{cl}, \mu_0[\mathcal{N}], \lambda[\mathcal{N}], \Omega[\mathcal{N}], \mathcal{N}] \Big|_{\mu_{\text{phys}}, \lambda_{\text{phys}}, \Omega_{\text{phys}} = \text{const}} \quad (19)$$

are finite and convergent in the limit $\mathcal{N} \rightarrow \infty$. In other words,

$$\lim_{\mathcal{N} \rightarrow \infty} \mathcal{N} \frac{d}{d\mathcal{N}} \left(\mathcal{Z}^N[\mathcal{N}] \Gamma_{m_1 n_1; \dots; m_N n_N}[\mu_0[\mathcal{N}], \lambda[\mathcal{N}], \Omega[\mathcal{N}], \mathcal{N}] \right) = 0 . \quad (20)$$

This implies the renormalisation group equation

$$\lim_{\mathcal{N} \rightarrow \infty} \left(\mathcal{N} \frac{\partial}{\partial \mathcal{N}} + N\gamma + \mu_0^2 \beta_{\mu_0} \frac{\partial}{\partial \mu_0^2} + \beta_{\lambda} \frac{\partial}{\partial \lambda} + \beta_{\Omega} \frac{\partial}{\partial \Omega} \right) \Gamma_{m_1 n_1; \dots; m_N n_N}[\mu_0, \lambda, \Omega, \mathcal{N}] = 0 , \quad (21)$$

where

$$\beta_{\mu_0} = \frac{1}{\mu_0^2} \mathcal{N} \frac{\partial}{\partial \mathcal{N}} \left(\mu_0^2[\mu_{\text{phys}}, \lambda_{\text{phys}}, \Omega_{\text{phys}}, \mathcal{N}] \right) , \quad (22)$$

$$\beta_{\lambda} = \mathcal{N} \frac{\partial}{\partial \mathcal{N}} \left(\lambda[\mu_{\text{phys}}, \lambda_{\text{phys}}, \Omega_{\text{phys}}, \mathcal{N}] \right) , \quad (23)$$

$$\beta_{\Omega} = \mathcal{N} \frac{\partial}{\partial \mathcal{N}} \left(\Omega[\mu_{\text{phys}}, \lambda_{\text{phys}}, \Omega_{\text{phys}}, \mathcal{N}] \right) , \quad (24)$$

$$\gamma = \mathcal{N} \frac{\partial}{\partial \mathcal{N}} \left(\ln \mathcal{Z}[\mu_{\text{phys}}, \lambda_{\text{phys}}, \Omega_{\text{phys}}, \mathcal{N}] \right) . \quad (25)$$

4 One-loop computations

Defining $(\Delta J)_{mn} := \sum_{p,q \in \mathbb{N}^2} \Delta_{mn;pq} J_{pq}$ we write (parts of) the generating functional of connected Green's functions up to second order in λ :

$$\begin{aligned}
W[J] = & \ln Z[0] + 4\pi^2 \theta^2 \sum_{m,n,k,l \in \mathbb{N}^2} \frac{1}{2} J_{mn} \Delta_{mn;kl} J_{kl} \\
& - (4\pi^2 \theta^2) \frac{\lambda}{4!} \sum_{m,n,k,l \in \mathbb{N}^2} \left\{ (\Delta J)_{ml} (\Delta J)_{lk} (\Delta J)_{kn} (\Delta J)_{nm} \right. \\
& + \frac{1}{4\pi^2 \theta^2} \left(\Delta_{nm;kn} (\Delta J)_{ml} (\Delta J)_{lk} + \Delta_{kn;lk} (\Delta J)_{nm} (\Delta J)_{ml} \right. \\
& \quad \left. \left. + \Delta_{nm;ml} (\Delta J)_{lk} (\Delta J)_{kn} + \Delta_{lk;ml} (\Delta J)_{kn} (\Delta J)_{nm} \right) \right. \\
& + \frac{1}{4\pi^2 \theta^2} \left(\Delta_{nm;lk} (\Delta J)_{kn} (\Delta J)_{ml} + \Delta_{kn;ml} (\Delta J)_{nm} (\Delta J)_{lk} \right) \\
& \left. + \frac{1}{(4\pi^2 \theta^2)^2} \left((\Delta_{nm;kn} \Delta_{lk;ml} + \Delta_{kn;lk} \Delta_{nm;ml}) + \Delta_{nm;lk} \Delta_{kn;ml} \right) \right\} \\
& + \frac{\lambda^2}{2(4!)^2} \sum_{m,n,k,l,r,s,t,u \in \mathbb{N}^2} \left\{ \left[\left(\Delta_{ml;sr} \Delta_{lk;ts} (\Delta J)_{kn} (\Delta J)_{nm} + \Delta_{ml;sr} \Delta_{kn;ts} (\Delta J)_{lk} (\Delta J)_{nm} \right. \right. \right. \\
& \quad \left. \left. + \Delta_{ml;sr} \Delta_{nm;ts} (\Delta J)_{lk} (\Delta J)_{kn} + \Delta_{lk;sr} \Delta_{ml;ts} (\Delta J)_{kn} (\Delta J)_{nm} \right. \right. \\
& \quad \left. \left. + \Delta_{lk;sr} \Delta_{kn;ts} (\Delta J)_{ml} (\Delta J)_{nm} + \Delta_{lk;sr} \Delta_{nm;ts} (\Delta J)_{ml} (\Delta J)_{kn} \right. \right. \\
& \quad \left. \left. + \Delta_{kn;sr} \Delta_{ml;ts} (\Delta J)_{lk} (\Delta J)_{nm} + \Delta_{kn;sr} \Delta_{lk;ts} (\Delta J)_{ml} (\Delta J)_{nm} \right. \right. \\
& \quad \left. \left. + \Delta_{kn;sr} \Delta_{nm;ts} (\Delta J)_{ml} (\Delta J)_{lk} + \Delta_{nm;sr} \Delta_{ml;ts} (\Delta J)_{lk} (\Delta J)_{kn} \right. \right. \\
& \quad \left. \left. + \Delta_{nm;sr} \Delta_{lk;ts} (\Delta J)_{ml} (\Delta J)_{kn} + \Delta_{nm;sr} \Delta_{kn;ts} (\Delta J)_{ml} (\Delta J)_{lk} \right) \right. \\
& \quad \left. \times (\Delta J)_{ru} (\Delta J)_{ut} \right. \\
& \quad \left. + 5 \text{ permutations of } ts, sr, ru, ut \right] \\
& + \text{1PI-contributions with } \leq 2 \text{ } J\text{'s} + \text{1PR-contributions} \left. \right\} + \mathcal{O}(\lambda^3). \quad (26)
\end{aligned}$$

In second order in λ we get a huge number of terms so that we display only the 1PI contribution with four J 's.

For the classical field (16) we get $\phi_{mn}^{\text{cl}} = \sum_{p,q \in \mathbb{N}^2} \Delta_{nm;pq} J_{pq} + \mathcal{O}(\lambda)$ so that

$$J_{pq} = \sum_{r,s \in \mathbb{N}^2} G_{qp;rs} \phi_{rs}^{\text{cl}} + \mathcal{O}(\lambda). \quad (27)$$

The remaining part not displayed in (27) removes the 1PR-contributions when passing to

$\Gamma[\phi^{cl}]$. We thus obtain

$$\Gamma[\phi^{cl}] = \Gamma[0] + 4\pi^2\theta^2 \sum_{m,n,k,l \in \mathbb{N}^2} \frac{1}{2} \left\{ G_{mn;kl} + \frac{\lambda}{6(4\pi^2\theta^2)} \left(\delta_{ml} \sum_{p \in \mathbb{N}^2} \Delta_{pn;kp} + \delta_{kn} \sum_{p \in \mathbb{N}^2} \Delta_{mp;pl} \right) \right. \quad (28a)$$

$$\left. + \frac{\lambda}{6(4\pi^2\theta^2)} \Delta_{ml;kn} + \mathcal{O}(\lambda^2) \right\} \phi_{mn}^{cl} \phi_{kl}^{cl} \quad (28b)$$

$$+ 4\pi^2\theta^2 \sum_{m,n,k,l,r,s,t,u \in \mathbb{N}^2} \frac{\lambda}{4!} \left\{ \delta_{nk} \delta_{lr} \delta_{st} \delta_{um} \right. \quad (28c)$$

$$\left. - \frac{\lambda}{2(4!)(4\pi^2\theta^2)} \left(\sum_{p,q \in \mathbb{N}^2} (4\Delta_{mp;qs} \Delta_{pl;tq} \delta_{kn} \delta_{ur} + 4\Delta_{kp;qs} \Delta_{pn;tq} \delta_{ml} \delta_{ur} + 4\Delta_{pl;rq} \Delta_{mp;qu} \delta_{nk} \delta_{st} + 4\Delta_{pn;rq} \Delta_{kp;qu} \delta_{ml} \delta_{st}) \right) \right. \quad (28d)$$

$$\left. + \sum_{p \in \mathbb{N}^2} (4\Delta_{ml;ps} \Delta_{kn;tp} \delta_{ur} + 4\Delta_{kn;ps} \Delta_{ml;tp} \delta_{ur} + 4\Delta_{mp;ts} \Delta_{pl;ru} \delta_{nk} + 4\Delta_{pl;ts} \Delta_{mp;ru} \delta_{nk} + 4\Delta_{kp;ts} \Delta_{pn;ru} \delta_{ml} + 4\Delta_{pn;ts} \Delta_{kp;ru} \delta_{ml} + 4\Delta_{ml;rp} \Delta_{kn;pu} \delta_{st} + 4\Delta_{kn;rp} \Delta_{ml;pu} \delta_{st}) \right. \quad (28e)$$

$$\left. + \sum_{p,q \in \mathbb{N}^2} (4\Delta_{pl;qs} \Delta_{mp;tq} \delta_{nk} \delta_{ur} + 4\Delta_{pn;qs} \Delta_{kp;tq} \delta_{ml} \delta_{ur} + 4\Delta_{kp;rq} \Delta_{pn;qu} \delta_{ml} \delta_{st} + 4\Delta_{mp;rq} \Delta_{pl;qu} \delta_{nk} \delta_{st}) \right. \quad (28f)$$

$$\left. + 4\Delta_{ml;ts} \Delta_{kn;ru} + 4\Delta_{kn;ts} \Delta_{ml;ru} \right) + \mathcal{O}(\lambda^2) \left\} \phi_{mn}^{cl} \phi_{kl}^{cl} \phi_{rs}^{cl} \phi_{tu}^{cl} \quad (28g)$$

$$+ \mathcal{O}(\lambda^2) .$$

Here, (28a) contains the contribution to the planar two-point function and (28b) the contribution to the non-planar two-point function. Next, (28c) and (28d) contribute to the planar four-point function, whereas (28e), (28f) and (28g) constitute three different types of non-planar four-point functions.

Introducing the cut-off $p^i, q^i \leq \mathcal{N}$ in the internal sums over $p, q \in \mathbb{N}^2$, we split the effective action according to [2] as follows into a relevant/marginal and an irrelevant piece ($\Gamma[0]$ can be ignored):

$$\Gamma[\phi^{cl}] \equiv \Gamma_{\text{rel/marg}}[\phi^{cl}] + \Gamma_{\text{irrel}}[\phi^{cl}] , \quad (29)$$

$$\Gamma_{\text{rel/marg}}[\phi^{cl}] = 4\pi^2\theta^2 \sum_{m,n,k,l \in \mathbb{N}^2} \frac{1}{2} \left\{ G_{mn;kl} + \frac{\lambda}{6(4\pi^2\theta^2)} \delta_{ml} \delta_{kn} \left(2 \sum_{p^1, p^2=0}^{\mathcal{N}} \Delta_{0 \ p^1; p^2 \ 0} \right. \right. \\ \left. \left. + (m^1+n^1+m^2+n^2) \sum_{p^1, p^2=0}^{\mathcal{N}} (\Delta_{1 \ p^1; p^2 \ 1} - \Delta_{0 \ p^1; p^2 \ 0}) \right) + \mathcal{O}(\lambda^2) \right\} \phi_{mn}^{cl} \phi_{kl}^{cl} \\ + 4\pi^2\theta^2 \sum_{m,n,k,l \in \mathbb{N}^2} \frac{\lambda}{4!} \left\{ 1 - \frac{\lambda}{3(4\pi^2\theta^2)} \sum_{p^1, p^2=0}^{\mathcal{N}} (\Delta_{0 \ p^1; p^2 \ 0})^2 + \mathcal{O}(\lambda^2) \right\} \phi_{mn}^{cl} \phi_{nk}^{cl} \phi_{kl}^{cl} \phi_{lm}^{cl} . \quad (30)$$

To the marginal four-point function and the relevant two-point function there contribute only the projections to planar graphs with vanishing external indices. The marginal two-point function is given by the next-to-leading term in the discrete Taylor expansion around vanishing external indices.

In a regime where $\lambda[\mathcal{N}]$ is so small that the perturbative expansion is valid in (30), the irrelevant part Γ_{irrel} can be completely ignored. Comparing (30) with the initial action according to (2),(7) and (8), we have $\Gamma_{\text{rel/marg}}[\mathcal{Z}\phi^{cl}] = S[\phi^{cl}; \mu_{\text{phys}}, \lambda_{\text{phys}}, \Omega_{\text{phys}}]$ with

$$\mathcal{Z} = 1 - \frac{\lambda}{192\pi^2\theta} \sum_{p^1, p^2=0}^{\mathcal{N}} (\Delta_{0 p^2; p^2 0}^{1 p^1, p^1 1} - \Delta_{0 p^2; p^2 0}^{0 p^1, p^1 0}) + \mathcal{O}(\lambda^2), \quad (31)$$

$$\begin{aligned} \mu_{\text{phys}}^2 &= \mu_0^2 \left(1 + \frac{\lambda}{12\pi^2\theta^2\mu_0^2} \sum_{p^1, p^2=0}^{\mathcal{N}} (2\Delta_{0 p^2; p^2 0}^{0 p^1, p^1 0} - \Delta_{0 p^2; p^2 0}^{1 p^1, p^1 1}) \right. \\ &\quad \left. - \frac{\lambda}{96\pi^2\theta} \sum_{p^1, p^2=0}^{\mathcal{N}} (\Delta_{0 p^2; p^2 0}^{1 p^1, p^1 1} - \Delta_{0 p^2; p^2 0}^{0 p^1, p^1 0}) + \mathcal{O}(\lambda^2) \right), \end{aligned} \quad (32)$$

$$\begin{aligned} \lambda_{\text{phys}} &= \lambda \left(1 - \frac{\lambda}{12\pi^2\theta^2} \sum_{p^1, p^2=0}^{\mathcal{N}} (\Delta_{0 p^2; p^2 0}^{0 p^1, p^1 0})^2 \right. \\ &\quad \left. - \frac{\lambda}{48\pi^2\theta} \sum_{p^1, p^2=0}^{\mathcal{N}} (\Delta_{0 p^2; p^2 0}^{1 p^1, p^1 1} - \Delta_{0 p^2; p^2 0}^{0 p^1, p^1 0}) + \mathcal{O}(\lambda^2) \right), \end{aligned} \quad (33)$$

$$\Omega_{\text{phys}} = \Omega \left(1 + \frac{\lambda(1-\Omega^2)}{192\pi^2\theta\Omega^2} \sum_{p^1, p^2=0}^{\mathcal{N}} (\Delta_{0 p^2; p^2 0}^{1 p^1, p^1 1} - \Delta_{0 p^2; p^2 0}^{0 p^1, p^1 0}) + \mathcal{O}(\lambda^2) \right). \quad (34)$$

Solving (32), (33) and (34) for the bare quantities, we obtain to one-loop order

$$\begin{aligned} &\mu_0^2[\mu_{\text{phys}}, \lambda_{\text{phys}}, \Omega_{\text{phys}}, \mathcal{N}] \\ &= \mu_{\text{phys}}^2 \left(1 - \frac{\lambda_{\text{phys}}}{12\pi^2\theta^2\mu_{\text{phys}}^2} \sum_{p^1, p^2=0}^{\mathcal{N}} \Delta_{0 p^2; p^2 0}^{0 p^1, p^1 0} \right. \\ &\quad \left. + \frac{\lambda_{\text{phys}}}{96\pi^2\theta} \left(1 + \frac{8}{\theta\mu_{\text{phys}}^2} \right) \sum_{p^1, p^2=0}^{\mathcal{N}} (\Delta_{0 p^2; p^2 0}^{1 p^1, p^1 1} - \Delta_{0 p^2; p^2 0}^{0 p^1, p^1 0}) + \mathcal{O}(\lambda_{\text{phys}}^2) \right), \end{aligned} \quad (35)$$

$$\begin{aligned} &\lambda[\mu_{\text{phys}}, \lambda_{\text{phys}}, \Omega_{\text{phys}}, \mathcal{N}] \\ &= \lambda_{\text{phys}} \left(1 + \frac{\lambda_{\text{phys}}}{12\pi^2\theta^2} \sum_{p^1, p^2=0}^{\mathcal{N}} (\Delta_{0 p^2; p^2 0}^{0 p^1, p^1 0})^2 + \frac{\lambda_{\text{phys}}}{48\pi^2\theta} \sum_{p^1, p^2=0}^{\mathcal{N}} (\Delta_{0 p^2; p^2 0}^{1 p^1, p^1 1} - \Delta_{0 p^2; p^2 0}^{0 p^1, p^1 0}) \right. \\ &\quad \left. + \mathcal{O}(\lambda_{\text{phys}}^2) \right), \end{aligned} \quad (36)$$

$$\begin{aligned}
& \Omega[\mu_{\text{phys}}, \lambda_{\text{phys}}, \Omega_{\text{phys}}, \mathcal{N}] \\
&= \Omega_{\text{phys}} \left(1 - \frac{\lambda_{\text{phys}}(1-\Omega_{\text{phys}}^2)}{192\pi^2\theta\Omega_{\text{phys}}^2} \sum_{p^1, p^2=0}^{\mathcal{N}} \left(\Delta_{\begin{smallmatrix} 1 & p^1 \\ 0 & p^2 \end{smallmatrix}; \begin{smallmatrix} p^1 & 1 \\ p^2 & 0 \end{smallmatrix}} - \Delta_{\begin{smallmatrix} 0 & p^1 \\ 0 & p^2 \end{smallmatrix}; \begin{smallmatrix} p^1 & 0 \\ p^2 & 0 \end{smallmatrix}} \right) + \mathcal{O}(\lambda_{\text{phys}}^2) \right). \quad (37)
\end{aligned}$$

Inserting (12) into (36) we can now compute the β_λ -function (23) up to one-loop order, omitting the index $_{\text{phys}}$ on μ^2 and Ω for simplicity:

$$\begin{aligned}
\beta_\lambda = & \frac{\lambda_{\text{phys}}^2}{48\pi^2} \mathcal{N} \frac{\partial}{\partial \mathcal{N}} \sum_{p^1, p^2=0}^{\mathcal{N}} \left\{ \left(\frac{{}_2F_1\left(1, \frac{\mu_0^2\theta}{8\Omega} - \frac{1}{2}(p^1+p^2) \middle| \frac{(1-\Omega)^2}{(1+\Omega)^2}\right)}{(1+\Omega)^2\left(1 + \frac{\mu_0^2\theta}{8\Omega} + \frac{1}{2}(p^1+p^2)\right)} \right)^2 \right. \\
& + \frac{p^1(1-\Omega)^2 {}_2F_1\left(3, \frac{1+\mu_0^2\theta}{8\Omega} - \frac{1}{2}(p^1+p^2+1) \middle| \frac{(1-\Omega)^2}{(1+\Omega)^2}\right)}{(1+\Omega)^4\left(\frac{1}{2} + \frac{\mu_0^2\theta}{8\Omega} + \frac{1}{2}(p^1+p^2)\right)\left(\frac{3}{2} + \frac{\mu_0^2\theta}{8\Omega} + \frac{1}{2}(p^1+p^2)\right)\left(\frac{5}{2} + \frac{\mu_0^2\theta}{8\Omega} + \frac{1}{2}(p^1+p^2)\right)} \\
& \left. + \frac{{}_2F_1\left(1, \frac{\mu_0^2\theta}{8\Omega} - \frac{1}{2}(p^1+p^2+1) \middle| \frac{(1-\Omega)^2}{(1+\Omega)^2}\right)}{2(1+\Omega)^2\left(\frac{3}{2} + \frac{\mu_0^2\theta}{8\Omega} + \frac{1}{2}(p^1+p^2)\right)} - \frac{{}_2F_1\left(1, \frac{\mu_0^2\theta}{8\Omega} - \frac{1}{2}(p^1+p^2) \middle| \frac{(1-\Omega)^2}{(1+\Omega)^2}\right)}{2(1+\Omega)^2\left(1 + \frac{\mu_0^2\theta}{8\Omega} + \frac{1}{2}(p^1+p^2)\right)} + \mathcal{O}(\lambda_{\text{phys}}) \right\}. \quad (38)
\end{aligned}$$

Symmetrising the numerator in the second line $p^1 \mapsto \frac{1}{2}(p^1+p^2)$ and using the expansions

$$\begin{aligned}
{}_2F_1\left(\begin{matrix} 1, & a-p \\ & b+p \end{matrix} \middle| z\right) &= \frac{1}{1+z} + \frac{z(a+b) + z^2(a+b-2)}{p(1+z)^3} + \mathcal{O}(p^{-2}), \\
{}_2F_1\left(\begin{matrix} 3, & a-p \\ & b+p \end{matrix} \middle| z\right) &= \frac{1}{(1+z)^3} + \mathcal{O}(p^{-1}), \quad (39)
\end{aligned}$$

which are valid for large p , we obtain up to irrelevant contributions vanishing in the limit $\mathcal{N} \rightarrow \infty$

$$\begin{aligned}
\beta_\lambda &= \frac{\lambda_{\text{phys}}^2}{48\pi^2} \mathcal{N} \frac{\partial}{\partial \mathcal{N}} \sum_{p^1, p^2=0}^{\mathcal{N}} \frac{1}{(1+\Omega_{\text{phys}}^2)^2} \frac{1}{(1+p^1+p^2)^2} \left\{ 1 + \frac{(1-\Omega_{\text{phys}}^2)^2}{2(1+\Omega_{\text{phys}}^2)} - \frac{(1+\Omega_{\text{phys}}^2)}{2} \right\} \\
&+ \mathcal{O}(\lambda_{\text{phys}}^3) + \mathcal{O}(\mathcal{N}^{-1}) \\
&= \frac{\lambda_{\text{phys}}^2}{48\pi^2} \frac{(1-\Omega_{\text{phys}}^2)}{(1+\Omega_{\text{phys}}^2)^3} + \mathcal{O}(\lambda_{\text{phys}}^3) + \mathcal{O}(\mathcal{N}^{-1}). \quad (40)
\end{aligned}$$

Similarly, one obtains

$$\beta_\Omega = \frac{\lambda_{\text{phys}}\Omega_{\text{phys}}}{96\pi^2} \frac{(1-\Omega_{\text{phys}}^2)}{(1+\Omega_{\text{phys}}^2)^3} + \mathcal{O}(\lambda_{\text{phys}}^2) + \mathcal{O}(\mathcal{N}^{-1}), \quad (41)$$

$$\begin{aligned}
\beta_{\mu_0} &= -\frac{\lambda_{\text{phys}}}{48\pi^2\theta\mu_{\text{phys}}^2(1+\Omega_{\text{phys}}^2)} \left(4\mathcal{N} \ln(2) + \frac{(8+\theta\mu_{\text{phys}}^2)\Omega_{\text{phys}}^2}{(1+\Omega_{\text{phys}}^2)^2} \right) \\
&+ \mathcal{O}(\lambda_{\text{phys}}^2) + \mathcal{O}(\mathcal{N}^{-1}), \quad (42)
\end{aligned}$$

$$\gamma = \frac{\lambda_{\text{phys}}}{96\pi^2} \frac{\Omega_{\text{phys}}^2}{(1+\Omega_{\text{phys}}^2)^3} + \mathcal{O}(\lambda_{\text{phys}}^2) + \mathcal{O}(\mathcal{N}^{-1}). \quad (43)$$

5 Discussion

We have computed the one-loop β - and γ -functions in real four-dimensional duality-covariant noncommutative ϕ^4 -theory. Remarkably, this model has a one-loop contribution to the wavefunction renormalisation which compensates partly the contribution from the planar one-loop four-point function to the β_λ -function. The one-loop β_λ -function is non-negative and vanishes in the distinguished case $\Omega = 1$ of the duality-invariant model, see (3). At $\Omega = 1$ also the β_Ω -function vanishes. This is of course expected (to all orders), because for $\Omega = 1$ the propagator (12) is diagonal, $\Delta_{\substack{m^1 & n^1 & k^1 & l^1 \\ m^2 & n^2 & k^2 & l^2}} \Big|_{\Omega=1} = \frac{\delta_{m^1 l^1} \delta_{k^1 n^1} \delta_{m^2 l^2} \delta_{k^2 n^2}}{\mu_0^2 + (4/\theta)(m^1 + m^2 + n^1 + n^2 + 2)}$, so that the Feynman graphs never generate terms with $|m^i - l^i| = |n^i - k^i| = 1$ in (8).

The similarity of the duality-invariant theory with the exactly solvable models discussed in [4] suggests that also the β_λ -function vanishes to all orders for $\Omega = 1$. The crucial differences between our model with $\Omega = 1$ and [4] is that we are using *real* fields, for which it is not so clear that the construction of [4] can be applied. But the planar graphs of a real and a complex ϕ^4 -model are very similar so that we expect identical β_λ -functions (possibly up to a global factor) for the complex and the real model. Since a main feature of [4] was the independence on the dimension of the space, the model with $\Omega = 1$ and matrix cut-off \mathcal{N} should be (more or less) equivalent to a two-dimensional model, which has a mass renormalisation only [8]. Therefore, we conjecture a vanishing β_λ -function in four-dimensional duality-invariant noncommutative ϕ^4 -theory to all orders.

The most surprising result is that the one-loop β_Ω -function also vanishes for $\Omega \rightarrow 0$. We cannot directly set $\Omega = 0$, because the hypergeometric functions in (38) become singular and the expansions (39) are not valid. Moreover, the power-counting theorems of [2], which we used to project to the relevant/marginal part of the effective action (30), also require $\Omega > 0$. However, in the same way as in the renormalisation of two-dimensional noncommutative ϕ^4 -theory [8], it is possible to switch off Ω very weakly with the cut-off \mathcal{N} , e.g. with

$$\Omega = e^{-\left(\ln(1+\ln(1+\mathcal{N}))\right)^2}. \quad (44)$$

The decay (44) for large \mathcal{N} over-compensates the growth of any polynomial in $\ln \mathcal{N}$, which according to [2] is the bound for the graphs contributing to a renormalisation of Ω . On the other hand, (44) does not modify the expansions (39). Thus, in the limit $\mathcal{N} \rightarrow \infty$, we have constructed the usual noncommutative ϕ^4 -theory given by $\Omega = 0$ in (2) at the one-loop level. It would be very interesting to know whether this construction of the noncommutative ϕ^4 -theory as the limit of a sequence (44) of duality-covariant ϕ^4 -models can be extended to higher loop order.

We also notice that the one-loop β_λ - and β_Ω -functions are independent of the noncommutativity scale θ . There is, however a contribution to the one-loop mass renormalisation via the dimensionless quantity $\mu_{\text{phys}}^2 \theta$, see (42).

Acknowledgement

We thank Helmut Neufeld for interesting discussions about the calculation of β -functions.

References

- [1] S. Minwalla, M. Van Raamsdonk and N. Seiberg, “Noncommutative perturbative dynamics,” *JHEP* **0002** (2000) 020 [arXiv:hep-th/9912072].
- [2] H. Grosse and R. Wulkenhaar, “Renormalisation of ϕ^4 theory on noncommutative \mathbb{R}^4 in the matrix base,” arXiv:hep-th/0401128.
- [3] E. Langmann and R. J. Szabo, “Duality in scalar field theory on noncommutative phase spaces,” *Phys. Lett. B* **533** (2002) 168 [arXiv:hep-th/0202039].
- [4] E. Langmann, R. J. Szabo and K. Zarembo, “Exact solution of quantum field theory on noncommutative phase spaces,” *JHEP* **0401** (2004) 017 [arXiv:hep-th/0308043].
- [5] K. G. Wilson and J. B. Kogut, “The Renormalization Group And The Epsilon Expansion,” *Phys. Rept.* **12** (1974) 75.
- [6] J. Polchinski, “Renormalization And Effective Lagrangians,” *Nucl. Phys. B* **231** (1984) 269.
- [7] H. Grosse and R. Wulkenhaar, “Power-counting theorem for non-local matrix models and renormalisation,” arXiv:hep-th/0305066.
- [8] H. Grosse and R. Wulkenhaar, “Renormalisation of ϕ^4 theory on noncommutative \mathbb{R}^2 in the matrix base,” *JHEP* **0312** (2003) 019 [arXiv:hep-th/0307017].