

# ON CERTAIN GRADED LIE ALGEBRAS ARISING IN NONCOMMUTATIVE GEOMETRY\*

R. MATTHES, G. RUDOLPH AND R. WULKENHAAR

Institut für Theoretische Physik der Universität Leipzig  
Augustusplatz 10/11, 04109 Leipzig, Germany

Given an algebra, a finite projective right module and a differential algebra over this algebra, a graded Lie algebra with derivation is constructed. It is shown that the algebraic structure of the Mainz-Marseille approach to the standard model may be obtained making use of this general construction in a special case. Thereby, a rigorous mathematical link between Connes' noncommutative geometry and the Mainz-Marseille approach is established.

PACS number: 02.40.-k

## 1. Introduction

The ideas of Connes, cf. [2] and [3], have been the starting point for numerous attempts to construct unified field theories using the tools of noncommutative geometry, the main achievement being, perhaps, the identification of the Higgs field as a gauge field. Slightly more modest seems to be – at first sight – the Mainz-Marseille approach, [4], which reaches essentially the same aims without using the precise geometrical notions of Connes, starting from a certain  $\mathbb{Z}_2$ -graded Lie algebra with derivation. We can, however, show that the algebraic structure of this latter approach can be derived in the scheme of Connes. The main point is to use a finite projective module (a notion which was avoided by the Mainz-Marseille group) and a differential algebra over the (underlying) algebra to construct a graded Lie algebra with derivation, which may be mapped by a partial homomorphism onto the graded Lie algebra of the Mainz-Marseille approach. Thereby we are able to give a precise geometrical meaning to all objects appearing there.

We will here only sketch the main mathematical ideas and refer to our paper [6] for all details. Moreover, we only mention, that there is a nice physical application of our method: Avoiding the “projection” to the Mainz-Marseille algebra, but nevertheless using the ideas of [4], it is possible to derive the standard model from the simplest two-point K-cycle originally used in [3] to derive the electroweak theory, see the last section of [6] and for details [7].

---

\*Presented at the II German–Polish Symposium “New Ideas in the Theory of Fundamental Interactions”, Zakopane, Poland, September 1995.

## 2. $\Omega_D A$ for the simplest two-point K-cycle

We will freely use several notions of noncommutative geometry whose definition seems to be by now standard. We refer to [2, 3, 5, 6] and the references given there for a detailed presentation. The first notion we will use is that of an even K-cycle  $(A, h, \pi, D, \Gamma)$  over an algebra. Every such K-cycle gives rise to a differential algebra  $\Omega_D A$  over  $A$ . We will need the following example:

Let  $X$  be a  $N = 2n$ -dimensional compact Riemannian spin manifold, let  $L^2(X, S)$  be the Hilbert space of square integrable sections of the spinor bundle, let  $D^{cl}$  be the classical Dirac operator, and let  $\gamma^{N+1}$  be the product of  $N$  orthonormal sections of the linear part of the Clifford bundle. Moreover, let

$$\mathcal{M} = \begin{pmatrix} 0 & M \\ M^* & 0 \end{pmatrix}, \quad \tilde{\Gamma} = \begin{pmatrix} 1_{n \times n} & 0 \\ 0 & -1_{n \times n} \end{pmatrix},$$

where  $M \in M_n \mathbb{C}$  with  $MM^* \notin \mathbb{C}1_{n \times n}$ . Consider the K-cycle consisting of

$$A = C^\infty(X) \otimes \mathbb{C}^2, \quad h = L^2(X, S) \otimes (\mathbb{C}^n \oplus \mathbb{C}^n) = L^2(X, S) \otimes \mathbb{C}^n \otimes \mathbb{C}^2,$$

$$D = D^{cl} \otimes id + \gamma^{N+1} \otimes \mathcal{M} = \begin{pmatrix} D^{cl} \otimes 1_{n \times n} & \gamma^{N+1} \otimes M \\ \gamma^{N+1} \otimes M^* & D^{cl} \otimes 1_{n \times n} \end{pmatrix},$$

$$\pi \begin{pmatrix} f_1 \\ f_2 \end{pmatrix} = \begin{pmatrix} f_1 \otimes 1_{n \times n} & 0 \\ 0 & f_2 \otimes 1_{n \times n} \end{pmatrix}, \quad \Gamma = \gamma^{N+1} \otimes \tilde{\Gamma},$$

This K-cycle was used in [3] for a construction of the Weinberg-Salam theory.

An explicit description of  $\Omega_D A$  for this K-cycle was given in [5]. The results obtained there may be summarized as follows: Let us denote  $M_1^t = (MM^*)^t$ ,  $M_2^t = M_1^t M$ ,  $M_4^t = (M^* M)^t$ ,  $M_3^t = M^* M_4^t$  and notice that there is a positive integer  $m$  such that in each series  $(M_q^t)_{t=0,1,\dots}$  ( $q = 1, 2, 3, 4$ ) just the first  $m$  terms are linearly independent in  $M_n \mathbb{C}$ . Then we have

$$\Omega_D^k A = \begin{pmatrix} \bigoplus_{t=0}^m \Lambda^{k-2t} \otimes \mathbb{C}M_1^t & \bigoplus_{t=0}^m \Lambda^{k-2t-1} \gamma^{N+1} \otimes \mathbb{C}M_2^t \\ \bigoplus_{t=0}^m \Lambda^{k-2t-1} \gamma^{N+1} \otimes \mathbb{C}M_3^t & \bigoplus_{t=0}^m \Lambda^{k-2t} \otimes \mathbb{C}M_4^t \end{pmatrix},$$

where  $\Lambda^k$  denotes the space of differential k-forms on  $X$  and the right multiplication with  $\gamma^{N+1}$  is nothing but (a certain variant of) the Hodge star. The product  $\bullet$  in  $\Omega_D A$  is given by

$$\begin{pmatrix} \alpha_1 \otimes M_1^{t_1} & \alpha_2 \gamma^{N+1} \otimes M_2^{t_2} \\ \alpha_3 \gamma^{N+1} \otimes M_3^{t_3} & \alpha_4 \otimes M_4^{t_4} \end{pmatrix} \bullet \begin{pmatrix} \beta_1^{l_1} \otimes M_1^{s_1} & \beta_2^{l_2} \gamma^{N+1} \otimes M_2^{s_2} \\ \beta_3^{l_3} \gamma^{N+1} \otimes M_3^{s_3} & \beta_4^{l_4} \otimes M_4^{s_4} \end{pmatrix} \\ = \begin{pmatrix} \alpha_1 \wedge \beta_1^{l_1} \otimes M_1^{t_1+s_2} & \alpha_1 \wedge \beta_2^{l_2} \gamma^{N+1} \otimes M_2^{t_1+s_2} \\ +(-1)^{l_3} \alpha_2 \wedge \beta_3^{l_3} \otimes M_1^{t_2+s_3+1} & +(-1)^{l_4} \alpha_2 \wedge \beta_4^{l_4} \otimes M_2^{t_2+s_4} \\ (-1)^{l_1} \alpha_3 \wedge \beta_1^{l_1} \gamma^{N+1} \otimes M_3^{t_3+s_1} & (-1)^{l_2} \alpha_3 \wedge \beta_2^{l_2} \otimes M_4^{t_3+s_2+1} \\ +\alpha_4 \wedge \beta_3^{l_3} \gamma^{N+1} \otimes M_3^{t_4+s_3} & +\alpha_4 \wedge \beta_4^{l_4} \otimes M_4^{t_4+s_4} \end{pmatrix}, \quad (1)$$

where we put  $M_i^t = 0$  for  $t > m$ , and the upper index of the  $\beta$ 's denotes the form degree. This is just multiplication of  $2 \times 2$ -matrices combined with exterior product of differential forms plus suitable signs arising from the exchange of a differential form with  $\gamma^{N+1}$ , and the following rules for the multiplication of the  $M_i^t$  (coming also from matrix multiplication):  $M_2^t M_3^s = M_1^{t+s+1}$ ,  $M_3^t M_2^s = M_4^{t+s+1}$ ,  $M_i^t M_j^s = M_{k(i,j)}^{t+s}$  for the other values of  $(i, j)$  ( $k(1, 1) = 1$ ,  $k(1, 2) = 2$ ,  $k(2, 4) = 2$ ,  $k(3, 1) = 3$ ,  $k(4, 3) = 3$ ,  $k(4, 4) = 4$ ). The differential  $\hat{d}$  can be written

$$\hat{d} = d + [\omega^{N+1}, \cdot]_g,$$

where  $d$  is the componentwise usual exterior differential, i.e.

$$d \begin{pmatrix} \alpha_1 \otimes M_1^{t_1} & \alpha_2 \gamma^{N+1} \otimes M_2^{t_2} \\ \alpha_3 \gamma^{N+1} \otimes M_3^{t_3} & \alpha_4 \otimes M_4^{t_4} \end{pmatrix} = \begin{pmatrix} d\alpha_1 \otimes M_1^{t_1} & d\alpha_2 \gamma^{N+1} \otimes M_2^{t_2} \\ d\alpha_3 \gamma^{N+1} \otimes M_3^{t_3} & d\alpha_4 \otimes M_4^{t_4} \end{pmatrix},$$

$\omega^{N+1} = -i \begin{pmatrix} 0 & \gamma^{N+1} \otimes M \\ \gamma^{N+1} \otimes M^* & 0 \end{pmatrix} \in \Omega_D^1 A$ , and  $[\cdot, \cdot]_g$  denotes the graded commutator with respect to the product  $\bullet$ . Notice that  $\Omega_D A$  is also a differential algebra with differential  $d$ .

### 3. The algebraic structure of the Mainz-Marseille approach

The central mathematical object of this approach is a certain  $\mathbb{Z}_2$ -graded Lie algebra with derivation contained in the  $\mathbb{Z}_2$ -graded differential algebra  $\Lambda(X) \otimes M_4\mathbb{C}$ , the latter one being considered as the  $\mathbb{Z}_2$ -graded tensor product of  $\Lambda(X)$  and  $M_4\mathbb{C}$ . Even and odd parts of  $M = \begin{pmatrix} A & B \\ C & D \end{pmatrix} \in M_4\mathbb{C}$  are defined to be  $M_0 = \begin{pmatrix} A & 0 \\ 0 & D \end{pmatrix}$  and  $M_1 = \begin{pmatrix} 0 & B \\ C & 0 \end{pmatrix}$ . With the usual matrix multiplication,  $M_4\mathbb{C}$  becomes a  $\mathbb{Z}_2$ -graded algebra, and with the corresponding graded commutator a  $\mathbb{Z}_2$ -graded Lie algebra, which we denote by  $pl(2, 2)$ . The graded differential is introduced as the graded commutator with the odd element  $\mathbf{m} = -i \begin{pmatrix} 0 & 1_{2 \times 2} \\ 1_{2 \times 2} & 0 \end{pmatrix}$ . It is also a graded differential of the graded Lie subalgebra  $sl(2, 2) = \{M \in M_4\mathbb{C} | tr(\Gamma_0 M) = 0\}$  of  $pl(2, 2)$ , where  $\Gamma_0 = \begin{pmatrix} 1_{2 \times 2} & 0 \\ 0 & -1_{2 \times 2} \end{pmatrix} \in M_4\mathbb{C}$ . Now, it is standard to define the graded tensor product  $\Lambda(X) \otimes M_4\mathbb{C}$  of differential algebras. Notice, in particular, that the differential  $\mathbf{d}$  can be written in the form  $\mathbf{d}(b = \beta \otimes M) = d\beta \otimes M + (-1)^{\partial\beta} \beta \otimes [\mathbf{m}, M] = db + [1 \otimes \mathbf{m}, b]_g$ . It turns out that  $\Lambda(X) \otimes spl(2, 2) \subset \Lambda(X) \otimes M_4\mathbb{C}$  as a graded differential Lie subalgebra. Now, define a graded Lie subalgebra of  $\Lambda(X) \otimes spl(2, 2)$  by

$$\Lambda(X) \otimes spl(2, 1) = \{b \in \Lambda(X) \otimes spl(2, 2) | b = \mathbf{e}b\mathbf{e}\},$$

where  $\mathbf{e} = \begin{pmatrix} 1_{2 \times 2} & 0 \\ 0 & \begin{pmatrix} 1 & 0 \\ 0 & 0 \end{pmatrix} \end{pmatrix}$ . Elements of  $\Lambda(X) \otimes spl(2, 1)$  just have zeroes in the last row and column. The differential  $\mathbf{d}$  descends to a derivation (not a differential!) of  $\Lambda(X) \otimes spl(2, 1)$  given by

$$\mathcal{D}b = \mathbf{e}db = db + [1 \otimes \mathbf{e}me, b]_g.$$

A connection in the Mainz-Marseille approach is an expression

$$\nabla = \mathbf{e}d + \mathbf{a}$$

with

$$\mathbf{a} = -\mathbf{a}^* = \begin{pmatrix} A_{11} & A_{12} & -i\Phi_1 & 0 \\ A_{21} & A_{22} & -i\Phi_2 & 0 \\ -i\Phi_1 & -i\Phi_2 & B & 0 \\ 0 & 0 & 0 & 0 \end{pmatrix} \in \Lambda(X) \otimes spl(2, 1)$$

and  $A_{ij} = -\bar{A}_{ji} \in \Lambda^1(X)$ ,  $B = -\bar{B} \in \Lambda^1(X)$ ,  $A_{11} + A_{22} = B$ ,  $\Phi_i \in \Lambda^0(X)$ . The curvature of such a connection is defined by  $\mathbf{f} = \nabla^2 = \mathbf{e}(\mathbf{d}\mathbf{e})(\mathbf{d}\mathbf{e})\mathbf{e} + \mathcal{D}\mathbf{a} + \frac{1}{2}[\mathbf{a}, \mathbf{a}]_g$ . Gauge transformations are defined on the infinitesimal level:  $\gamma_t(\mathbf{a}) =$

$$\mathbf{a} - \mathcal{D}t + [t, \mathbf{a}]_g \text{ with } t = -t^* = \begin{pmatrix} T_{11} & T_{12} & 0 & 0 \\ T_{21} & T_{22} & 0 & 0 \\ 0 & 0 & T_{33} & 0 \\ 0 & 0 & 0 & 0 \end{pmatrix} \in \Lambda(X) \otimes spl(2, 1), \text{ where}$$

$T_{ij} = -\bar{T}_{ji} \in \Lambda^0(X)$  and  $tr(\Gamma_0 t) = 0$ . In [4], the gauge and Higgs bosons of the electroweak theory were unified in the ‘‘connection form’’  $\mathbf{a}$ . Notice that the above constructions may be easily generalized using  $M_{2p}\mathbb{C}$  instead of  $M_4\mathbb{C}$ , which leads to  $pl(p, p)$ ,  $spl(p, p)$  etc.

#### 4. A general construction of graded Lie algebras with derivation

Let us start with the following data: Let  $A$  be a unital  $*$ -algebra over  $\mathbb{C}$ , let  $(\Lambda_A, \bullet, *, d)$  be an involutive differential algebra over  $A$  ( $\Lambda_A^0 = A$ ), and let  $\mathcal{E} = eA^p$  be a finite projective right  $A$ -module with Hermitian structure  $(\cdot, \cdot)_{\mathcal{E}}$ . We put

$$\mathcal{E}^* = \bigoplus_{k=0}^{\infty} \mathcal{E}^k, \text{ where } \mathcal{E}^k = \mathcal{E} \otimes_A \Lambda_A^k.$$

$\mathcal{E}^*$  is a right  $\Lambda_A$ -module in a natural way, and there are natural extensions of the Hermitian metric to mappings  $(\cdot, \cdot)_{\mathcal{E}}^{k,l} : \mathcal{E}^k \times \mathcal{E}^l \longrightarrow \Lambda_A^{k+l}$ . Now, we define

$$\mathcal{H} = \bigoplus_{k=0}^{\infty} \mathcal{H}^k, \quad \mathcal{H}^k = Hom_A(\mathcal{E}, \mathcal{E}^k).$$

$\mathcal{H}$  can be given the structure of an associative  $\mathbb{N}$ -graded involutive algebra over  $\mathbb{C}$ : The product  $\bullet$  is defined by

$$(\rho^k \bullet \rho^l)(\xi) = (id_{\mathcal{E}} \otimes_A \bullet) \circ (\rho^k \otimes_A id_{\Lambda_A^l}) \circ \rho^l(\xi).$$

$id_{\mathcal{E}}$  is the unit for this multiplication, and the involution is defined by

$$(\xi, (\rho^k)^*(\xi^l))_{\mathcal{E}}^{0,k} = (\rho^k(\xi), \xi^l)_{\mathcal{E}}^{k,0}.$$

With the graded commutator,  $\mathcal{H}$  becomes also an  $\mathbb{N}$ -graded Lie algebra, and it acts from the left on  $\mathcal{E}^*$ :  $\rho^k \bullet \xi^l = (id_{\mathcal{E}} \otimes_A \bullet) \circ (\rho^k \otimes_A id_{\Lambda_A^l})(\xi^l)$ . Finally, there is a graded derivation  $D_{\mathcal{H}} : \mathcal{H}^k \rightarrow \mathcal{H}^{k+1}$  inherited from the canonical compatible connection  $\nabla_0$  on  $\mathcal{E}$ , which stems from the differential  $d$  of  $\Lambda_A$ :

$$(D_{\mathcal{H}}\rho^k)(\xi) = \nabla_0(\rho^k(\xi)) - (-1)^k \rho^k \bullet (\nabla_0(\xi)).$$

$D_{\mathcal{H}}$  fails to be a differential:

$$D_{\mathcal{H}}^2(\rho) = \Theta_0 \bullet \rho - \rho \bullet \Theta_0,$$

where  $\Theta_0$  is the curvature of  $\nabla_0$ . For the curvature of a connection  $\nabla = \nabla_0 + \rho$ , one obtains

$$\Theta = \Theta_0 + \mathcal{D}_{\mathcal{H}}\rho + \rho \bullet \rho.$$

These definitions have a nice matrix form: Let  $(\epsilon_i)_{i=1}^p$  be the canonical basis of  $A^p$  ( $\epsilon_i$  having the unit of  $A$  as entry at the  $i$ -th place, zeroes at the other places). Then, the projection  $e$  is given by  $e(\epsilon_i) = \epsilon_j e_{ji}$ ,  $e_{ji} \in A$  with  $e_{ij}e_{jk} = e_{ik}$ .  $\xi \in \mathcal{E}$  is characterized by  $(e\xi)_i = e_{ij}\xi_j = \xi_i$ . An element  $\rho \in \mathcal{H}^k$  is characterized by a matrix  $(\rho_{ij})_{i,j=1}^p$ ,  $\rho_{ij} \in \Lambda_A^k$ , with  $e_{ij}\rho_{jk}e_{kl} = \rho_{il}$  (in short,  $e\rho e = \rho$ ), the multiplication in  $\mathcal{H}$  is given by matrix multiplication,  $(\rho \bullet \rho')_{ij} = \rho_{ik} \bullet \rho'_{kj}$ , and the derivation  $D_{\mathcal{H}}$  is given by componentwise action of  $d$ :  $(D_{\mathcal{H}}\rho)_{ij} = e_{ik}d\rho_{kl}e_{lj}$  ( $D_{\mathcal{H}}\rho = ed\rho e$ ). Notice that  $(\Theta_0)_{ij} = e_{ik}de_{kl} \bullet de_{lm}e_{mj}$ .

In order to come from these general definition of an algebra with derivation to the algebraic structure of the Mainz-Marseille approach, we have first to specify the data of our definition. For the chosen case, it is possible to introduce a suitable condition of tracefreeness on  $\mathcal{H}$  and a certain surjective mapping whose application just leads to the structures of the foregoing section. First, we take the algebra  $A = C^\infty(X) \otimes \mathbb{C}^2$  and the differential algebra  $\Lambda_A = \Omega_D A$  of the K-cycle described in the example of section 2. For this case, and for any module  $\mathcal{E}$ , we can construct a certain graded Lie subalgebra  $\mathcal{H}_0$  of  $\mathcal{H}$  as follows: We define a  $\mathbb{C}$ -linear map  $T_\lambda : \Lambda_A \rightarrow \Lambda(X)$  by

$$T_\lambda \left( \left( \begin{array}{cc} \alpha_1 \otimes M_1^{t_1} & \alpha_2 \gamma^{N+1} \otimes M_2^{t_2} \\ \alpha_3 \gamma^{N+1} \otimes M_3^{t_3} & \alpha_4 \otimes M_4^{t_4} \end{array} \right) \right) = \alpha_1 + \alpha_4.$$

This is a generalized trace in the sense that  $T_\Lambda(\Gamma_\Lambda[\lambda, \lambda]_g) = 0$ , where  $\Gamma_\Lambda = \begin{pmatrix} 1 \otimes 1_{n \times n} & 0 \\ 0 & -1 \otimes 1_{n \times n} \end{pmatrix} \in A$ . Now, we define  $T_{\mathcal{H}} : \mathcal{H} \rightarrow \Lambda(X)$  by

$$T_{\mathcal{H}}(\rho) = \sum_{i=0}^p T_\Lambda(\Gamma_\Lambda \rho_{ii}),$$

which is also a generalized trace:  $T_{\mathcal{H}}([\rho, \rho']_g) = 0$ . Therefore,  $\mathcal{H}_0 = \bigoplus_{k=0}^{\infty} \mathcal{H}_0^k$ ,  $\mathcal{H}_0^k = \{\rho \in \mathcal{H}^k | T_{\mathcal{H}}(\rho) = 0\}$ , is a graded Lie subalgebra of  $\mathcal{H}$ .

Recall that there are two differentials  $\hat{d}$  and  $d$  on  $\Omega_D A$ . For both we can construct, using the corresponding compatible connections  $\hat{\nabla}_0$  and  $\nabla_0$ , graded derivations  $\hat{\mathcal{D}}_{\mathcal{H}}$  and  $\mathcal{D}_{\mathcal{H}}$  of  $\mathcal{H}$ , which turn out to be also graded derivations of  $\mathcal{H}_0$ . They are related by

$$\hat{\mathcal{D}}_{\mathcal{H}}\rho = \mathcal{D}_{\mathcal{H}}\rho + [\mu, \rho]_g,$$

where  $\mu = e(1_{p \times p} \otimes \omega^{N+1})e \in \mathcal{H}^1$ .

To come to the Mainz Marseille setting, we now have to perform two steps:

1. In matrix representation, elements of  $\mathcal{H}$  are  $p \times p$ -matrices with entries from  $\Lambda_A \subset \Lambda(X) \otimes \text{End}(\mathbb{C}^n) \otimes M_2\mathbb{C}$ . We treat them now as  $2 \times 2$ -matrices with entries from  $\Lambda(X) \otimes \text{End}(\mathbb{C}^n) \otimes M_p\mathbb{C}$ . This is just going from one standard representation of a Kronecker product of matrices to the other one. Moreover, we can remove the  $\gamma^{N+1}$  without losing information. Thus, we get an injection

$$i : \Lambda_A \otimes M_p\mathbb{C} \rightarrow \Lambda(X) \otimes M_p\mathbb{C} \otimes \text{End}(\mathbb{C}^n) \otimes M_2\mathbb{C}$$

of vector spaces. Elements of  $i(\Lambda_A^k \otimes M_p\mathbb{C})$  have the form

$$\begin{pmatrix} A_1^{k-2t_1} \otimes M_1^{t_1} & A_2^{k-2t_2-1} \otimes M_2^{t_2} \\ A_3^{k-2t_3-1} \otimes M_3^{t_3} & A_4^{k-2t_4} \otimes M_4^{t_4} \end{pmatrix}$$

with  $A_q^l \in \Lambda^l(X) \otimes M_p\mathbb{C}$ . Moreover, we have  $i(e) = \begin{pmatrix} e_1 & 0 \\ 0 & e_4 \end{pmatrix}$  with  $e_q = e_q^2 = e_q^* \in C^\infty(X) \otimes M_p\mathbb{C}$ . Elements of  $i(\mathcal{H})$  are characterized by  $A_1 = e_1 A_1 e_1$ ,  $A_2 = e_1 A_2 e_4$ ,  $A_3 = e_4 A_3 e_1$ ,  $A_4 = e_4 A_4 e_4$ , those of  $i(\mathcal{H}_0)$  in addition by  $\text{tr} A_1 = \text{tr} A_4$ . Transporting the product  $\bullet$  of  $\mathcal{H}$  leads to a product of the same form as in  $\Lambda_A = \Omega_D A$ , formula (1). One has to replace there  $\alpha \rightarrow A$ ,  $\beta \rightarrow B$ , one has to omit  $\gamma^{N+1}$  and one has to interpret  $\wedge$  as exterior product of forms combined with multiplication of  $p \times p$ -matrices.

2. We define a surjection  $\mathbf{p} : i(\Lambda_A \otimes M_p\mathbb{C}) \rightarrow \Lambda(X) \otimes M_{2p}\mathbb{C}$  by

$$\mathbf{p} \begin{pmatrix} A_1 \otimes M_1^{t_1} & A_2 \otimes M_2^{t_2} \\ A_3 \otimes M_3^{t_3} & A_4 \otimes M_4^{t_4} \end{pmatrix} = \begin{pmatrix} A_1 & A_2 \\ A_3 & A_4 \end{pmatrix}.$$

**Theorem 1**

- (i)  $\mathbf{p} \circ i(\mathcal{H}_0) = \{b \in \Lambda(X) \otimes \text{spl}(p, p) | b = ebe\}$   
with  $\mathbf{e} = i(e) = \begin{pmatrix} e_1 & 0 \\ 0 & e_4 \end{pmatrix}$  (see above).
- (ii)  $(\mathbf{p} \circ i(\rho))^* = \mathbf{p} \circ i(\rho^*)$ ,  $\rho \in \mathcal{H}$ .
- (iii)  $\mathbf{p} \circ i([\rho^k, \rho^l]_g) = [\mathbf{p} \circ i(\rho^k), \mathbf{p} \circ i(\rho^l)]_g$  for  $k + l \leq 2m + 1$ ,  $\rho^k \in \mathcal{H}^k$ ,  $\rho^l \in \mathcal{H}^l$ .
- (iv)  $\mathbf{p} \circ i(\hat{\mathcal{D}}_{\mathcal{H}}(\rho^k)) = \mathcal{D}(\mathbf{p} \circ i(\rho^k))$  for  $k \leq 2m$ ,  $\rho^k \in \mathcal{H}^k$ .

Notice that also the analogue  $\mathbf{p} \circ i(\mathcal{D}_{\mathcal{H}}(\rho^k)) = d(\mathbf{p} \circ i(\rho^k))$  of (iv) is true.

The theorem says that  $\mathbf{p} \circ i$  is a partial homomorphism of  $\mathbb{Z}_2$ -graded involutive Lie algebras with derivation. This mapping is not injective on  $\mathcal{H}$  or  $\mathcal{H}_0$ , its restriction, however, to any sum of subsequent homogeneous components  $(\Lambda_A^k \oplus \Lambda_A^{k+1}) \otimes M_p \mathbb{C}$  is injective. Since we assume  $MM^* \notin \mathbb{C}1$ , we have  $m \geq 1$ . Therefore, in particular,  $\mathbf{p}$  is a monomorphism on the graded Lie subalgebra  $i(\mathcal{H}^0 \oplus \mathcal{H}^1)$ , and it commutes with the derivation of elements of  $i(\mathcal{H}^0)$  and  $i(\mathcal{H}^1)$ . However, under the application of  $\mathbf{p}$  the  $\mathbb{N}$ -grading of  $i(\mathcal{H})$  is lost and only a  $\mathbb{Z}_2$ -grading remains. It is now easy to see, that we arrive at the algebraic setting of the Mainz-Marseille approach starting with the choice  $p = 2$  and

$$e = \begin{pmatrix} \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix} & 0 \\ 0 & \begin{pmatrix} 1 & 0 \\ 0 & 0 \end{pmatrix} \end{pmatrix}. \text{ Using the above theorem, it is almost obvious}$$

that under the mapping  $\mathbf{p} \circ i$  the geometric objects living in the projective module  $\mathcal{E} = eA^2$  are transformed into corresponding objects of the Mainz-Marseille scheme. In particular, due to the partial injectivity of  $\mathbf{p} \circ i$  discussed above, no information about the objects relevant for gauge theories (connections and curvatures) is lost. Moreover, the scheme is completed by giving a natural definition of the (noninfinitesimal) gauge group and of the module where the connection acts.

**References**

- [1] A. Connes, *Essay on Physics and Non-commutative Geometry*, preprint IHES/M/89/69.
- [2] A. Connes, *Non commutative geometry*, Academic Press, New York 1994.
- [3] A. Connes and J. Lott, *The Metric Aspect of Noncommutative Geometry*, Proceedings of 1991 Cargese summer conference, ed. by J. Fröhlich et al. (Plenum, New York 1992).
- [4] R. Coquereaux, G. Esposito-Farèse and F. Scheck, *Noncommutative Geometry and Graded Algebras in Electroweak Interactions*, Int. J. Mod. Phys. A **7** (1992) 6555–6593.
- [5] R. Matthes, G. Rudolph and R. Wulkenhaar, *On the structure of a differential algebra used by Connes and Lott*, Rep. Math. Phys. **38** (1996) 45-66.

- [6] R. Matthes, G. Rudolph and R. Wulkenhaar, *On a certain construction of graded Lie algebras with derivation*, J. Geom. Phys. **20** (1996) 107-141.
- [7] R. Wulkenhaar, *Deriving the standard model from the simplest two-point K-cycle*, J. Math. Phys. **37** (1996) 3797-3814.