

Bounded Martin's Maximum and strong cardinals

Ralf Schindler¹

*Institut für Mathematische Logik und Grundlagenforschung
Westfälische Wilhelms-Universität Münster, Einsteinstr. 62, 48149 Münster, Germany*

rds@math.uni-muenster.de

<http://wwwmath.uni-muenster.de/math/inst/logik/org/staff/rds/>

Abstract

We show that if Bounded Martin's Maximum (BMM) holds then for every $X \in V$ there is an inner model with a strong cardinal containing X . We also discuss various open questions which are related to BMM.

1 Introduction and statement of the result.

This paper strengthens one of the results of [6]. Bounded Martin's Maximum (BMM, for short) is the statement that whenever $\mathbb{P} \in V$ is a stationary set preserving forcing notion then

$$H_{\omega_2}^V \prec_{\Sigma_1} H_{\omega_2}^{V^{\mathbb{P}}}.$$

Bounded forcing axioms were introduced in [2] (as weakenings of the “unbounded” forcing axioms PFA and MM). Todorćević showed (cf. [8]) that BMM implies that $2^{\aleph_0} = \aleph_2$. (This was later improved by J. Moore who showed that already the Bounded Proper Forcing Axiom implies $2^{\aleph_0} = \aleph_2$; cf. [4].) We refer the reader to [9, Section 10.3] for a discussion of BMM. We proved in [6, Theorem 1.3] that if BMM holds then for every $X \in V$, $X^\#$ exists. The purpose of the present note is to prove the following.

Theorem 1.1 *Suppose that BMM holds. Then for every $X \in V$ there is an inner model with a strong cardinal containing X .*

We do not know if Theorem 1.1 gives the optimal lower bound for the consistency strength of BMM. Woodin has shown in unpublished work (cf. [10]) that $\omega + 1$ many Woodin cardinals are an upper bound. We refer the reader to [6] for further background information.

¹The main result of this note was proven in February 2004 while the author was a guest at the CRM in Barcelona. He would like to thank Neus Portet and Joan Bagaria for their warm hospitality.

Whereas [6] constructs, assuming $\text{BMM} + V$ is not closed under the $\#$ operator, a strictly decreasing sequence of functions from ω_1 to ω_1 , the key new idea here is to construct, assuming $\text{BMM}+$ there is some set X which is not in an inner model with a strong cardinal, a strictly decreasing sequence of functions from ω_1 to the set of all countable mice, where “decreasing” means “decreasing in the mouse order.”

2 The proof.

Let us assume that BMM holds thruout this section. We shall prove that there is an inner model with a strong cardinal. The reader will gladly verify that the argument to follow “relativizes” to any $X \in V$, yielding a proof of Theorem 1.1.

Let us suppose that there is no inner model with a strong cardinal. We may thus let K denote the core model (cf. [3]). If \mathcal{M} is a premouse and $\alpha \leq \mathcal{M} \cap \text{OR}$ then we say that α is overlapped in \mathcal{M} just in case there is some extender $E_\nu^{\mathcal{M}}$ such that $\text{crit}(E_\nu^{\mathcal{M}}) < \alpha$ and $\nu \geq \alpha$. As there is no inner model with a strong cardinal, in K there is no E_ν^K such that $\text{crit}(E_\nu^K)$ is overlapped in K .

We shall use the following notation. Let $\mathcal{M} = J_\alpha[E]$ be a premouse, and let $A \subset \bar{\alpha}$ for some $\bar{\alpha} < \alpha$. Then by $\mathcal{M}[A]$ we denote the transitive set (structure) $J_\alpha[E, A]$. Of course, in general $\mathcal{M}[A]$ will not be a premouse (or not even be a model of a reasonable fragment of ZFC). Nevertheless, models of the form $\mathcal{M}[A]$ will play a key rôle in what follows.

By \leq^* we shall denote the pre-well-ordering of mice (cf. for instance [7]). I.e., if \mathcal{M} and \mathcal{N} are mice and \mathcal{T}, \mathcal{U} is the coiteration of \mathcal{M}, \mathcal{N} then $\mathcal{M} \leq^* \mathcal{N}$ if and only if $\mathcal{M}_\infty^{\mathcal{T}} \sqsubseteq \mathcal{M}_\infty^{\mathcal{U}}$ (in which case $[0, \infty)_{\mathcal{T}}$ contains no drop). We shall write $\mathcal{M} <^* \mathcal{N}$ iff $\mathcal{M} \leq^* \mathcal{N}$ and $\neg(\mathcal{N} \leq^* \mathcal{M})$.

Using the Dodd-Jensen Lemma and the fact that there are no degenerate iterations of mice, one can show that \leq^* is indeed a pre-well-ordering. The following Lemma will thus give the desired contradiction.

Lemma 2.1 (*BMM+ there is no inner model with a strong cardinal.*) *There is a sequence $\mathcal{S} = (A_n, C_n, (\mathcal{N}_{n,\alpha} : \alpha \in C_n) : n < \omega)$ such that for every $n < \omega$, $A_n \subset \omega_1$, C_n is a club subset of ω_1 , $C_n \subset C_{n-1}$ if $n > 0$, and for every $\alpha \in C_n$, $\mathcal{N}_{n,\alpha}$ is a sound mouse with $\rho_\omega(\mathcal{N}_{n,\alpha}) = \kappa \geq \alpha$, where κ is the largest cardinal of $\mathcal{N}_{n,\alpha}$, and κ is not overlapped in $\mathcal{N}_{n,\alpha}$, $(A_n \cap \alpha)_{\text{odd}}$ codes the mouse $\mathcal{N}_{n,\alpha} \upharpoonright \kappa$,² and $\mathcal{N}_{n,\alpha}[A_n \cap \alpha] \models$*

²If A is a set of ordinals then by A_{odd} we mean the set $\{\alpha : 2\alpha + 1 \in A\}$. The coding here and in what follows is to be understood as being according to some standard soft coding device. For instance, if M is transitive and of size ζ then there is some $E \subset \zeta^2$ and some isomorphism $\sigma : (\zeta; E) \cong (M; \in)$; via Gödel’s pairing function, E may be construed as a subset of ζ which codes M .

“ α is countable”; moreover, $\mathcal{N}_{n,\alpha} <^* \mathcal{N}_{n-1,\alpha}$ if $n > 0$ and $\alpha \in C_n$.

The proof of Lemma 2.1 exploits a version of the “faster reshaping forcing” which we had introduced in [6].

Let $n \in \omega$, and let us assume that $\mathcal{S} \upharpoonright n$ has been constructed. We aim to construct A_n , C_n , and $(\mathcal{N}_{n,\alpha} : \alpha \in C_n)$.

By BMM, it suffices to force the existence of these objects with the desired properties by a stationary set preserving forcing notion.

By [5], there is a σ -closed forcing notion \mathbb{P} such that in $V^{\mathbb{P}}$, CH holds, there is some $A^+ \subset \omega_1$ with $H_{\omega_2} = K \parallel_{\omega_2}[A^+]$, and $K \parallel_{\omega_2}$ has a largest cardinal κ which is not overlapped in $K \parallel_{\omega_2}$. We may assume that A^+ is chosen such that $H_{\omega_1} = K \parallel_{\omega_1}[A^+]$.

Let us for the rest of this argument assume that $n > 0$. The case $n = 0$ is an easier variant of what is to come. We thus may and shall assume that A_{odd}^+ is the join of A_{n-1} and a code of $K \parallel_{\kappa}$. Let $C \subset C_{n-1}$ be club and such that there is a sequence $(\mathcal{M}_\alpha : \alpha \in C)$ of mice such that for every $\alpha \in C$, $(A^+ \cap \alpha)_{\text{odd}}$ codes the join of $A_{n-1} \cap \alpha$ and a code of \mathcal{M}_α . Let κ_α denote the height of \mathcal{M}_α for $\alpha \in C$. The following will be crucial.

Claim 1. Let $\pi : \bar{K}[A^+ \cap \alpha] \cong X \prec (K \parallel_{\omega_2}[A^+]; \in, A^+)$, where X is countable and $\alpha = X \cap \omega_1 \in C$. Then $\mathcal{M}_\alpha \triangleleft \bar{K}$, $\kappa_\alpha = \pi^{-1}(\kappa)$ is the largest cardinal of \bar{K} , and $\bar{K} \leq^* \mathcal{N}_{n-1,\alpha}$.

PROOF OF CLAIM 1. Everything except for $\bar{K} \leq^* \mathcal{N}_{n-1,\alpha}$ easily follows by the elementarity of π . Let us write $\mathcal{N} = \mathcal{N}_{n-1,\alpha}$. Suppose Claim 1 to be false; i.e., $\mathcal{N} <^* \bar{K}$. We shall argue that $\bar{K}[A^+ \cap \alpha] \models$ “ α is countable,” which is of course nonsense.

Because \bar{K} does not have any active extenders with indices between $\pi^{-1}(\kappa)$ and its height, it is easy to see that there must be now some $\xi < \bar{K} \cap \text{OR}$ such that $\mathcal{N} <^* \bar{K} \parallel \xi$. Let \mathcal{T}, \mathcal{U} denote the coiteration of $\mathcal{N}, \bar{K} \parallel \xi$. We know that $[0, \infty)_{\mathcal{T}}$ contains no drops, and that $\mathcal{M}_\infty^{\mathcal{T}} \trianglelefteq \mathcal{M}_\infty^{\mathcal{U}}$.

Let $\rho_\omega(\mathcal{N}) = \eta$. Because $(A_{n-1} \cap \alpha)_{\text{odd}}$ codes $\mathcal{N} \parallel \eta$, $\mathcal{N} \parallel \eta \in L[A_{n-1} \cap \alpha] \subset L[A^+ \cap \alpha]$.

Let $\bar{\mathcal{T}}, \bar{\mathcal{U}}$ denote the coiteration of $\mathcal{N} \parallel \eta, \bar{K} \parallel \xi$. Clearly, $\bar{\mathcal{T}}$ is an “initial segment” of \mathcal{T} , and $\bar{\mathcal{U}}$ is an “initial segment” of \mathcal{U} . Moreover, $\mathcal{N} \parallel \eta <^* \bar{K} \parallel \xi$, so that $[0, \infty)_{\bar{\mathcal{T}}}$ contains no drops and $\mathcal{M}_\infty^{\bar{\mathcal{T}}} \trianglelefteq \mathcal{M}_\infty^{\bar{\mathcal{U}}}$. If we had $\text{lh}(\mathcal{T}) > \text{lh}(\bar{\mathcal{T}})$ then, as η is not overlapped in \mathcal{N} and $\rho_\omega(\mathcal{N}) = \eta$, $\mathcal{M}_\infty^{\mathcal{T}}$ would be non-sound, although $\mathcal{N} <^* \bar{K}$. Therefore, $\text{lh}(\mathcal{T}) = \text{lh}(\bar{\mathcal{T}})$. But then we must also have that $\text{lh}(\mathcal{U}) = \text{lh}(\bar{\mathcal{U}})$, as η is the largest cardinal of \mathcal{N} .

Because $\mathcal{N} \parallel \eta \in L[A^+ \cap \alpha]$ and $\bar{K} \parallel \xi \in L[\bar{K} \parallel \xi]$, we know that $\pi_{0\infty}^{\bar{\mathcal{T}}}$ as well as $\mathcal{M}_\infty^{\bar{\mathcal{U}}} = \mathcal{M}_\infty^{\mathcal{U}}$ are elements of $L[A^+ \cap \alpha, \bar{K} \parallel \xi]$.

As $\mathcal{M}_\infty^T \trianglelefteq \mathcal{M}_\infty^u$, we thus have that $\mathcal{M}_\infty^T \in L[A^+ \cap \alpha, \bar{K} \parallel \xi]$. But

$$\mathcal{N} \cong h^{\mathcal{M}_\infty^T}(\text{ran}(\pi_{0\infty}^T) \cup \{\pi_{0\infty}^T(p_{\mathcal{N}})\}),$$

where $p_{\mathcal{N}}$ is the standard parameter of \mathcal{N} and $h^{\mathcal{M}_\infty^T}$ is an appropriate Skolem hull operator. This yields that in fact $\mathcal{N} \in L[A^+ \cap \alpha, \bar{K} \parallel \xi]$.

Now let $a \in {}^\omega\omega \cap \mathcal{N}[A_n \cap \alpha]$ code a well-order of ω of order-type α . We have shown that $a \in L[A^+ \cap \alpha, \bar{K} \parallel \xi]$. By [6], $V^{\mathbb{P}}$ is closed under the $\#$ operator. We therefore have that $(A^+ \cap \alpha, \bar{K} \parallel \xi)^\#$ exists and $a \in (A^+ \cap \alpha, \bar{K} \parallel \xi)^\#$. But $A^+ \cap \alpha$ and $\bar{K} \parallel \xi$ are both elements of $\bar{K}[A^+ \cap \alpha]$. Due to π , $\bar{K}[A^+ \cap \alpha]$ is closed under the $\#$ operator, so that in fact $a \in \bar{K}[A^+ \cap \alpha]$. But then α is countable in $\bar{K}[A^+ \cap \alpha]$, a contradiction. \square (Claim 1)

In $V^{\mathbb{P}}$, we shall now consider the following forcing notion, denoted by \mathbb{Q} . We let $(c, p) \in \mathbb{Q}$ if and only if c is a countable closed subset of C , $p: \max(c) \rightarrow 2$, and for every $\alpha \in c$ there is some sound mouse $\mathcal{N} \triangleq \mathcal{M}_\alpha$ such that κ_α is the largest cardinal of \mathcal{N} , κ_α is not overlapped in \mathcal{N} , $\rho_\omega(\mathcal{N}) = \kappa_\alpha$, $\mathcal{N}[A^+ \cap \alpha, p \upharpoonright \alpha] \models$ “ α is countable,” and $\mathcal{N} <^* \mathcal{N}_{n-1, \alpha}$. A condition (c', q) is stronger than (c, p) iff $\max(c') \geq \max(c)$, $c' \cap (\max(c) + 1) = c$, and $q \upharpoonright \max(c) = p$.

Claim 2. (“Extendability Lemma”) If $(c, p) \in \mathbb{Q}$ and $\alpha < \omega_1$ then there is some $(c', q) \leq (c, p)$ such that $\max(c') \geq \alpha$.

PROOF OF CLAIM 2. Given (c, p) and α , let us assume w.l.o.g. that $\alpha \geq \max(c) + \omega$, $\alpha \in C$, and $\alpha = X \cap \omega_1$ for some $X \prec K \parallel \omega_2[A^+]$, so that $X \cong \bar{K}$ for some $\bar{K} \triangleright \mathcal{M}_\alpha$ and $\bar{K} \leq^* \mathcal{N}_{n-1, \alpha}$ by Claim 1. Pick a code $x \in {}^\omega\omega$ for the ordinal α , and let $\text{dom}(q) = \alpha$, where $q \upharpoonright \text{dom}(p) = p$, $q(\max(c) + n) = x(n)$ for all $n < \omega$, and $q(\gamma) = 0$ for all $\gamma \geq \text{dom}(c) + \omega$. There will be some \mathcal{P} with $\bar{K} \parallel \pi^{-1}(\kappa) \triangleleft \mathcal{P} \triangleleft \bar{K}$ such that $\rho_\omega(\mathcal{P}) = \pi^{-1}(\kappa)$, $\pi^{-1}(\kappa)$ is the largest cardinal of \mathcal{P} , and $\mathcal{P}[A^+ \cap \alpha, q] \models$ “ α is countable.” Therefore, $(c \cup \{\alpha\}, q)$ is as desired. \square (Claim 2)

In order to finish the proof of Lemma 2.1 (and hence of Theorem 1.1) it now obviously suffices to verify the following.

Claim 3. \mathbb{Q} is stationary set preserving.

PROOF OF CLAIM 3. Let $S \subset \omega_1$ be stationary, and let $(c, p) \Vdash$ “ \dot{C} is a club subset of $\check{\omega}_1$.” We aim to construct some $(c', q) \leq (c, p)$ such that $(c', q) \Vdash$ “ $\dot{C} \cap \check{S} \neq \emptyset$.”

Let $X \prec (H_{\omega_2}; \in, A^+, \mathbb{Q}, (c, p))$ be transitive and of size \aleph_1 , and let $(X_i: i \leq \omega_1)$ be a continuous chain of countable elementary submodels of X approaching X (i.e., $X = X_{\omega_1}$). Recall that $H_{\omega_2} = K \upharpoonright_{\omega_2}[A^+]$ (in $V^{\mathbb{P}}$). Let

$$\pi: \bar{K}[A^+ \cap \alpha] \cong Y \prec (H_{\omega_2}; \in, A^+, \mathbb{Q}, (c, p), (X_i: i \leq \omega_1)),$$

where Y is countable and $\alpha = Y \cap \omega_1 = \omega_1^{\bar{H}[A^+ \cap \alpha]} < \omega_1$. We also may and shall assume that $\alpha \in S$. Let us write $\alpha_i = X_i \cap \omega_1 < \omega_1$ for $i < \omega_1$. Of course $(\alpha_i: i < \alpha) \in \bar{K}[A^+ \cap \alpha]$, where $(\alpha_i: i < \alpha)$ is cofinal in α . Let us pick (externally, i.e., in $V^{\mathbb{P}}$) a sequence $(i_n: n < \omega)$ which is cofinal in α ; hence $(\alpha_{i_n}: n < \omega)$ is cofinal in α as well. As $H_{\omega_1} = K \upharpoonright_{\omega_1}[A^+]$, $(H_{\omega_1})^{\bar{K}[A^+ \cap \alpha]} \subset \bigcup_{i < \alpha} X_i$, so that we may assume that in fact $(c, p) \in X_{i_0}$.

We shall now recursively construct sequences $((c_n, p_n): n < \omega)$ and $((c'_n, p'_n): n < \omega)$ of conditions in \mathbb{Q} with the following properties.

- (1) $(c_0, p_0) = (c, p)$,
- (2) $(c'_n, p'_n) \leq (c_n, p_n)$, and $(c'_n, p'_n) \in X_{i_{n+1}}$,
- (3) for all $\xi \in (\text{dom}(p'_n) \setminus \text{dom}(p_n)) \cap \{\alpha_i: i < \alpha\}$, $p'_n(\xi) = 1$ if and only if $\xi = \alpha_{i_n}$,
- (4) $(c_{n+1}, p_{n+1}) \leq (c'_n, p'_n)$, and $(c_{n+1}, p_{n+1}) \in X_{i_{n+1}}$,
- (5) there is some $\beta > \alpha_{i_n}$ such that $(c_{n+1}, p_{n+1}) \Vdash \check{\beta} \in \dot{C}$, and
- (6) $\max(c_{n+1}) > \alpha_{i_n}$.

In the light of (the proof of) Claim 2, there is no problem with this recursion.

Let us now set $c^* = \bigcup_{n < \omega} c_n \cup \{\alpha\}$ and $p^* = \bigcup_{n < \omega} p_n$. We're done if we can show that (c^*, p^*) is a condition, because then $(c^*, p^*) \Vdash \check{\alpha} \in \dot{S} \cap \dot{C}$.

Well, we have that $\bar{K}[A^+ \cap \alpha, p^*] \models \text{"}\alpha \text{ is countable,}"$ because for $\xi \in \{\alpha_i: i_0 \leq i < \alpha\}$, $p^*(\xi) = 1$ if and only if $\xi = \alpha_{i_n}$ for some $n < \omega$, so that $(\alpha_{i_n}: n < \omega) \in \bar{K}[A^+ \cap \alpha, p^*]$.

By Claim 1, $\bar{K} \triangleright \mathcal{M}_\alpha = \bar{K} \upharpoonright_{\pi^{-1}(\kappa)}$. Moreover, there will again certainly be some \mathcal{P} with $\bar{K} \upharpoonright_{\pi^{-1}(\kappa)} \triangleleft \mathcal{P} \triangleleft \bar{K}$ such that $\rho_\omega(\mathcal{P}) = \pi^{-1}(\kappa)$, $\pi^{-1}(\kappa)$ is the largest cardinal of \mathcal{P} , and $\mathcal{P}[A^+ \cap \alpha, p^*] \models \text{"}\alpha \text{ is countable.}"$ By Claim 1, $\mathcal{P} <^* \mathcal{N}_{n-1, \alpha}$, so that (c^*, p^*) is really a condition. □ (Claim 3)

□ (Lemma 2.1, Theorem 1.1)

3 Some problems.

Let $(f_\alpha: \alpha < \omega_2)$ denote “the” sequence of canonical functions from ω_1 to ω_1 . I.e., $f_\alpha(\nu) = \text{otp } g_\alpha \upharpoonright \nu$, where $g_\alpha: \omega_1 \rightarrow \alpha$ is bijective. The Club Bounding Principle, CBP for short, says that for every $f: \omega_1 \rightarrow \omega_1$ there is some $\alpha < \omega_2$ such that $f < f_\alpha$ on a club, i.e., such that $\{\nu: f(\nu) < f_\alpha(\nu)\}$ contains a club subset of ω_1 .

P. Larson has shown that CBP implies $P_2(\omega_1)^+$ (cf. [1, Fact 3.4]; cf. [1, Definition 3.1 (b)] on the definition of $P_2(\omega_1)^+$). On the other hand one has:

Lemma 3.1 *If BMM and $P_2(\omega_1)^+$ hold then CBP holds.*

PROOF SKETCH. Let $f: \omega_1 \rightarrow \omega_1$. If $P_2(\omega_1)^+$ holds then the natural forcing \mathbb{P}_f for adding a canonical function above f is easily seen to be stationary set preserving. \mathbb{P}_f is just the collection of all (c, p) such that c is a closed countable subset of ω_1 and $p: \max(c) \rightarrow \omega_2$ is such that for all $\nu \in c$, $f(\nu) < \text{otp } p''\nu$. \square (Lemma 3.1)

We do not know if BMM alone implies the Club Bounding Principle. In order to answer this question we need a better understanding of how to construct models of BMM. The same remark applies to the following questions.

Woodin introduced the principle ψ_{AC} and showed that ψ_{AC} implies that $2^{\aleph_0} = \aleph_2$. (Cf. [9, Definition 5.12 and Lemma 5.15].) We do not know if BMM implies that ψ_{AC} holds (we know that BMM implies that $2^{\aleph_0} = \aleph_2$, though, cf. [8]). But we have:

Lemma 3.2 *If BMM and $P_1(\omega_1)^+$ hold then ψ_{AC} holds.*

PROOF SKETCH. Cf. [1, Definition 3.1 (d)] on the definition of $P_1(\omega_1)^+$. Let $S, T \subset \omega_1$ both be stationary and costationary. By $P_1(\omega_1)^+$, the canonical forcing $\mathbb{P}_{S,T}$ for adding a witness to ψ_{AC} is easily seen to be stationary set preserving. $\mathbb{P}_{S,T}$ consists of all (c, p) such that c is a closed countable subset of ω_1 , $p: \max(c) \rightarrow \omega_2$, and for all $\nu \in c$, $\text{otp } p''\nu \in T \iff \nu \in S$. \square (Lemma 3.2)

It is not known if ψ_{AC} implies $P_1(\omega_1)^+$. By [1, Fact 3.1], ψ_{AC} implies CBP.

Another interesting question is whether BMM implies that $u_2 = \omega_2$. This question makes sense by [6]. If BMM holds then every real has a $\#$, and hence we may ask if u_2 , the second uniform indiscernible, is equal to ω_2 .

References

- [1] Aspero, D., and Welch, Ph., *Bounded Martin's Maximum, weak Erdős cardinals, and ψ_{AC}* , Journal Symb. Logic **67** (2002), pp. 1141-1152.
- [2] Goldstern, M., and Shelah, S., *The bounded proper forcing axiom*, Journal Symb. Logic **60** (1995), pp. 58-73.
- [3] Jensen, R., *The core model for non-overlapping extender sequences*, handwritten notes, Oxford.

- [4] Moore, J., *Set mapping reflection*, preprint.
- [5] Schindler, R., *Coding into K by reasonable forcing*, Trans. Amer. Math. Soc. **353** (2000), pp. 479-489.
- [6] Schindler, R., *Semi-proper forcing, remarkable cardinals, and Bounded Martin's Maximum*, Mathematical Logic Quarterly, submitted.
- [7] Steel, J., and Welch, Ph., Σ_3^1 *absoluteness and the second uniform indiscernible*, Israel Journal of Mathematics **104** (1998), pp. 157-190.
- [8] Todorcevic, S., *Generic absoluteness and the continuum*, Mathematical Research Letters **9** (2002), pp. 1-7.
- [9] Woodin, H., *The axiom of determinacy, forcing axioms, and the nonstationary ideal*, de Gruyter 1999.
- [10] Woodin, H., private communication, Aug. 03.